

**QUANTUM HAMILTON - JACOBI SOLUTION FOR
SPECTRA OF SEVERAL ONE DIMENSIONAL POTENTIALS
WITH SPECIAL PROPERTIES**

A thesis submitted in partial fulfilment of the requirements
for the award of the degree of

DOCTOR OF PHILOSOPHY

in

PHYSICS

by

S. SREE RANJANI



**SCHOOL OF PHYSICS
UNIVERSITY OF HYDERABAD
HYDERABAD - 500046, INDIA
JUNE 2004**

Declaration

I, ***S. Sree Ranjani***, hereby declare that the work reported in this dissertation titled, ***Quantum Hamilton - Jacobi solution for spectra of several one dimensional potentials with special properties***, is entirely original and has been carried out by me, under the supervision of ***Prof. A. K. Kapoor***, Department of Physics, School of Physics, University of Hyderabad.

To the best of my knowledge, no part of this dissertation was submitted for any degree of any other institute or university.

Place: Hyderabad

Date:

S Sree Ranjani

Certificate

This is to certify that the report entitled *Quantum Hamilton - Jacobi solution for spectra of several one dimensional potentials with special properties*, being submitted by *S Sree Ranjani*, bearing *Regd. No. 99phph02*, in partial fulfillment of the requirements for the award of *Doctor of Philosophy in Physics* by *University of Hyderabad*, is a bonafide work carried out at the *University of Hyderabad* under my supervision. The matter embodied in this report has not been submitted to any other institute or university for the award of any degree.

Dean,
School of Physics,
University of Hyderabad,
Hyderabad - 500046.

Prof. A. K. Kapoor,
School of Physics,
University of Hyderabad,
Hyderabad - 500046.

Dedicated to

My parents and my supervisor

Acknowledgments

*Catching the fleeting moment of a sunset,
feeling the soft caress of the wind,
the world I live for, the love I'd die for,
and all because, **you pass this way but once...***

I am glad, I came this way and I consider it a blessing to have worked with my supervisor Prof. A. K. Kapoor. The most beautiful human being! His humility, his patience and confidence are always my inspiration. His depth of understanding of the subject and his urge to teach has always left me amazed. There are not enough words to express my gratitude for his encouragement, understanding and support throughout. There were many times, when I kept working only because he believed in me, even when I lacked faith in myself. “Dear sir, I take this opportunity to thank you for everything and it was inspiring to work under you”.

I convey my sincere gratitude to Dr. P. K. Panigrahi for taking interest in my work and giving me constant guidance. Discussions with him were always lively and motivating. His valuable suggestions allowed me to explore different aspects of my work and in a way moulded it.

I thank the Dean, School of Physics, Prof. V. S. S. Sastry and the former dean, Prof. S. N. Kaul for providing congenial atmosphere for research.

I thank Prof. V. Srinivasan and Prof. S. Chaturvedi for the fascinating and enjoyable lectures. Though my interaction with them is very little, their presence is felt and revered. I thank Dr. M. Siva kumar, Dr. B. A. Bambah, Prof. C. Mukku and Dr. Anantha Lakshmi for their encouragement and support.

I am thankful to all of the faculty members of the School of Physics for the interesting courses and lectures in my M. Sc. These have helped me to enhance my understanding in Physics and nurtured my dream of doing research. I also thank all the non - teaching staff for their support, especially I thank Mr. K. Srinivas (Gaddam) for his help with my latex files and Mr. T. Abraham for his timely and cheerful help.

The long association with the campus has left me with a treasure trove of memories and experiences. The campus, with its blooming trees, lovely lakes, beautiful sunsets and skies, has always enchanted and mesmerized me. My life has become richer with the friends, I made in the campus. It's a pleasure to extend my heartfelt appreciation to my friends, Swati, Chandu, Kuki, Sarita, Charan, Sunil, Meri, Jo, Latha, Manoj, Madhu, Deepu, Nandu and many more, because of whom my days were filled with fun and laughter. The most memorable were the excited discussions, which flamed and sustained our zeal for physics. I acknowledge Mr. Seshagiri Rao for his encouragement. I thank my colleagues and friends, Phani, Ajith, Mani,

Raghu, Sathya, Rajneesh, Rizwan, Chaitanya, Sudarshan, Farzana, Kumar and many others for providing a friendly atmosphere and extending their help, when needed. I also acknowledge my seniors, Siddique, Gurappa, Hari Kumar, Sunil, Solomon, Raishma, Anita, Ramana, Phani Kumar, Naga Srinivas, Prem Kiran, Geojo and Sandhya for their encouragement and help. I am grateful to Charan and Rajneesh for the time and energy, they have spent on going through the manuscripts of my papers.

A special thanks to valli for always being there, when ever I needed. I thank her for believing in me and dreaming for me. A note of thanks to her family for the warmth and affection bestowed on me.

I thank my parents, in - laws, sister, grand parents and all other family members for their encouragement and support. I am grateful to them and to my friends for standing by me during the bad phase of ill health. I thank Mr. Bhaskar Rao, Mr. and Mrs. Dr. Ram Kumar for their timely help.

Last but not the least, I thank my husband, Ravi, for his unfailing love and affection for constantly and patiently encouraging me to push back my limits. Especially, I am deeply indebted to him and my mother for taking care of me and helping me to build my battered health and self esteem. Most of all for keeping my spirit alive.

*Two roads diverged in a wood, and
I took the one less travelled by,
and that has made all the difference.
- Robert Frost*

Contents

1	Introduction	1
2	Quantum Hamilton - Jacobi Formalism	12
2.1	Classical Hamilton - Jacobi theory	12
2.1.1	Action - angle variables	16
2.2	Quantum Hamilton - Jacobi Formalism	18
2.2.1	Singularities of the QMF	19
2.3	Hydrogen atom	23
2.3.1	Definition of p_c	25
2.3.2	Residue at the fixed pole $r = 0$	27
2.3.3	Calculation of energy eigenvalues and eigenfunctions	28
2.4	Change of variables in QHJ Formalism	30
3	Study of potentials exhibiting different spectra	36
3.1	Introduction	36
3.2	Phases of Supersymmetry	37
3.2.1	Form of QMF (χ)	38

3.2.2	Definition of the Classical momentum function	39
3.2.3	Residues at the fixed poles $y = \pm 1$	41
3.2.4	Case 1 : $(A - B) > 0$, $(A + B) > 0$	43
3.2.5	Case 2 : $(A - B) > 0$, $(A + B) < 0$	44
3.3	Phases of a periodic potential	45
3.3.1	Form of the QMF χ	48
3.3.2	Choice of the residues	50
3.3.3	Case 1 : Band spectrum	51
3.3.4	Case 2 : Bound state spectrum	54
4	Exactly Solvable Periodic Potentials	56
4.1	Introduction	56
4.2	General Lamé potential	59
4.2.1	Form of QMF χ	60
4.2.2	Explicit forms of the wave functions	64
4.3	Lamé potential with $j = 2$	65
4.4	Associated Lamé potential	69
4.4.1	Form of QMF (χ)	69
4.5	Associated Lamé potential with $j = 1$	71
5	Periodic Quasi Exactly Solvable Models	80
5.1	Introduction	80
5.2	QES condition and the forms of the wave functions	82
5.2.1	Form of QMF (χ)	83

5.2.2	Forms of wave function	84
5.3	Associated Lamé potential with a, b as integers	88
5.4	Associated Lamé potential with a and b as half integers	91
6	PT symmetric Hamiltonians	95
6.1	Introduction	95
6.2	Khare - Mandal model	96
6.2.1	Form of QMF (χ)	97
6.2.2	Forms of the wave function	99
6.2.3	Case 1 : $M = 3$	100
6.2.4	Case 2 : $M=2$	101
6.3	Complex Scarf - II potential	102
6.3.1	Case 1 : $ B > A + \frac{1}{4}$	105
6.3.2	Case 2 : $ B \leq A + \frac{1}{4}$	106
7	Conclusions	109

Chapter 1

Introduction

It is well known that there are many formulations of classical mechanics, like the Lagrangian, Hamiltonian, Poisson bracket and Hamilton - Jacobi formulation [1, 2]. Among these, the first two are the most important and most used formulations. The Hamilton - Jacobi formalism is an alternate formulation, which is not widely used and is more esoteric. Although, all these provide the same solutions to a given problem in an independent way, one needs to know about them because they provide different insights into the physics of the problem. Secondly, solving a particular problem might be easy in one formalism compared to the other. Hence, depending on the problem, one needs to choose a method, in which it can be solved easily. Similar to classical mechanics, there are various formulations of quantum mechanics [3]. Of these, some have been developed independently and some have their origins in classical mechanics. The quantum transformation theory along with the quantum Hamilton - Jacobi (QHJ) formalism, which is the subject of this thesis, belongs to the later group. The transformation theory

has been developed analogous to the classical theory, the roots of which, lie in the works of Dirac [4, 5], Jordan [6] and Schwinger [7]. The QHJ formalism, in its present form, has been proposed by Leacock and Padgett [8, 9], on the lines of its classical counterpart, to study stationary states.

In classical mechanics, the theory of canonical transformation and the Hamilton - Jacobi (HJ) theory are very well established [1, 2]. These provided powerful formal techniques, to deal with dynamical systems, which led to a better understanding of the Hamiltonian formalism and the phase space. The HJ theory provides an efficient tool to study the systems describing periodic motion. Making use of the action - angle variables, one can find the frequency of the periodic motion, without first finding the full solution of the equation of motion explicitly. Similarly, in the QHJ formalism, one can find the energy eigenvalues without solving the Schrödinger equation for the stationary states.

Analogous to the classical theory, the quantum momentum function (QMF) p is defined as dS/dx , where S is analogous to the Hamilton's characteristic function and will be called quantum action function. S satisfies the equation

$$\left(\frac{dS}{dx}\right)^2 - i\hbar \left(\frac{d^2S}{dx^2}\right) = 2m(E - V(x)), \quad (1.1)$$

which in the limit $\hbar \rightarrow 0$ goes to the classical Hamilton - Jacobi equation [1].

The solution of the Schrödinger equation, ψ is related to S by

$$\psi = \exp\left(\frac{iS}{\hbar}\right) \quad (1.2)$$

and one can see that p is the logarithmic derivative of ψ ,

$$p = -i\hbar \frac{d}{dx}(\ln \psi). \quad (1.3)$$

In the QHJ formalism, the quantum momentum function plays a central role. Here, unlike the conventional wave mechanics, where one has to solve the Schrödinger equation, one makes use of the quantum Hamilton - Jacobi equation for the QMF

$$p^2 - i\hbar \frac{dp}{dx} = 2m(E - V(x)), \quad (1.4)$$

which is a Riccati type equation. The method makes use of the singularity structure of the QMF to compute the energy eigenvalues in a simple and elegant fashion [8, 9]. Thus, for quantum mechanical systems, which allow periodic motion, one can obtain the bound state energy eigenvalues by defining the quantum action J , which is a generalization of the classical action variable and J satisfies the exact quantization condition,

$$J = J(E) = \frac{1}{2\pi} \oint_C p dx = n\hbar, \quad (1.5)$$

where C is the contour in the complex plane enclosing the n poles of the QMF on the real line. This quantization condition is exact and is a result of the oscillation theorem [10, 11] which states that the n^{th} excited state wave function has n zeros. From (1.3), one can see that these n zeros correspond to the n poles of the QMF and are called moving poles, as they are dependent on the initial conditions. It can be shown from (1.4) that these poles have a

residue $-i\hbar$. Thus, applying the Cauchy residue theorem to the integral in (4.1), one obtains the above quantization condition.

Along with these moving poles, the QMF will have other singularities originating from the potential which are called fixed singularities. Here, one makes a simplifying assumption that *the point at infinity is an isolated singular point*. This assumption is equivalent to saying that the QMF has finite number of moving poles in the complex domain. Evaluating the integral in (4.1), in terms of these other singular points and using the Cauchy residue theorem, one can obtain the energy eigenvalue expression. Due to the quadratic nature of the QHJ equation, all the residues at the fixed poles will have two values. Hence, one needs to choose the value of the residue, which will give the right physical behaviour. For this, one needs to define the QMF completely. The QMF is related to the classical momentum function p_c by the relation

$$\lim_{\hbar \rightarrow 0} p \rightarrow p_c, \quad (1.6)$$

which doubles up as a boundary condition for the QMF, as well as being the correspondence principle. Note that p_c is double valued and has to be defined precisely. The details of its definition are given in the next chapter.

Bhalla *et.al.* [12 - 14] used this formalism to compute the bound state spectra of one dimensional exactly solvable (ES) supersymmetric potentials [15, 16]. They have also demonstrated the applicability of the QHJ formalism to the potentials, which exhibit both broken and unbroken phases of SUSY [17, 14], depending on the range of the potential parameters. For a given po-

tential, it was shown that implementing the boundary conditions correctly, every value of the residue, at a given fixed pole was acceptable in a particular range of parameters and one can obtain the energy eigenvalues for both the phases of SUSY through the QHJ formalism. Extending this investigation to complicated potentials like Ginocchio, Natanzon etc, Jayanthi *et. al.* obtained the energy eigenvalues for these potentials and also obtained the wave functions for a few states [18].

After successfully obtaining the solutions for bound state energies, it was natural to look at the other potential models like the quasi - exactly solvable (QES) models [19]. These are termed as QES, because one cannot analytically obtain their entire energy spectrum but it is possible to obtain a few states analytically, when the potential parameters satisfy a specific condition known as the quasi - exact solvability condition. Applying QHJ formalism to these models it was found that unlike the case of ES models, where one obtains the energy expressions, one is lead to the quasi - exact solvability condition by the application of the quantization condition[20, 21]. In this case, the property of square integrability was used to choose the right value of residue instead of the boundary condition (1.6), due to the difficulty in its implementation to these models.

In the above investigations except in [18], the main emphasis was on obtaining the energy eigenvalues and not much attention was paid to eigenfunctions. We have shown in [22], that one can easily obtain the bound state eigenfunctions for one dimensional exactly solvable models [15] along with

the energy eigenstates through the QHJ formalism. In addition to these, the eigenfunctions and eigenvalues for the QES models, discussed in [20], were obtained in [21].

Having solved one dimensional ES and QES models, we have taken up the application of this formalism to more complicated problems like the periodic potentials [23 - 26] and the PT symmetric potentials [27, 28, 29]. The application of QHJ formalism to these models is not a straight forward extension of the method used for the bound state problems in one dimension. In particular, attention has to be paid to the choice of residues because square integrability condition cannot be insisted. Also the quantization condition holds as long as the oscillation theorem holds. For PT symmetric potentials, it is not clear whether a generalization of the oscillation theorem can be found. So we had to find an alternative, which will give the energy values, for which, the details are given in the later chapters.

The plan of the thesis is as follows. In the next chapter, we present an introduction to the QHJ formalism. We will describe in detail, how one can employ the QHJ formalism to obtain the energy eigenfunctions and eigenvalues, by explicitly working out the well known hydrogen atom potential. In chapter III, the study of Scarf [30] and Scarf - I [15, 13] potentials is taken up. Both the potentials are interesting models because they exhibit different spectra depending on the range of the potential parameters. The Scarf potential exhibits band structure for one parameter range and ordinary bound state spectrum for another range. The other potential Scarf - I exhibits the

broken and unbroken phases of SUSY in different regions of its potential parameters. For these two models, both the regions of the parameters and the corresponding energy spectra emerge naturally through the QHJ formalism.

The next two chapters deal with periodic potentials belonging to the family of elliptic potentials [23 - 26], namely the Lamé potential,

$$V(x) = j(j+1)m \operatorname{sn}^2(x, m) \quad (1.7)$$

with j being an integer and the associated Lamé potential (ALP) [26]

$$V(x) = a(a+1)m \operatorname{sn}^2(x, m) + b(b+1)m \frac{\operatorname{cn}^2(x, m)}{\operatorname{dn}^2(x, m)}. \quad (1.8)$$

Here, the functions $\operatorname{sn}(x, m)$, $\operatorname{cn}(x, m)$ and $\operatorname{dn}(x, m)$ are the elliptic functions with m being the elliptic modulus. The ALP is ES if $a = b = j$, j being an integer and is QES, when $a \neq b$ with $a + b, a - b$ being integers. Here, we show, how one can obtain the band edge solutions of these potentials using the QHJ formalism. This way of obtaining the solutions, turns out to be much simpler compared to the existing methods [24] and the comparison of the singularity structure with the ordinary ES and QES models is taken up [31,32].

Chapter IV deals with ES periodic potentials *i.e.* the Lamé and the ES case of the ALP. For these potentials, we obtain the general forms of the band edge wave functions in terms of the potential parameters. Explicit expressions for the energy and wave functions are worked out for specific examples [31]. In chapter V, we study the QES periodic potentials *i.e.* the ALP, when $a \neq b$ and a, b taking non - integer values. Here, we obtain the

quasi exact solvability conditions along with the general forms of the wave functions [32].

Chapter VI deals with the study of PT symmetric ES and QES solvable potentials. In this chapter, the complex Scarf potential and the Khare - Mandal model are discussed [33]. The comparison of the singularity structure of the QMF of these potentials with the ordinary ES and QES potential models is given. Chapter VII contains the concluding remarks along with a brief discussion on further applications of this work.

References

1. H. Goldstein, *Classical Mechanics* (Addison-Wesley/Narosa, Indian student edition, 1995)
2. E. J. Saletan and A. H. Cromer *Theoretical Mechanics* (John Wiley, New York, 1971).
3. D. F. Styer, *et. al. Am. J. Phys.* **70** (3) 288 (2002)
4. P. A. M. Dirac, *The Principles of Quantum Mechanics* (Oxford University Press, London, 1958).
5. P. A. M. Dirac, *Rev. Mod. Phys.* **17**, 195 (1945) ; P. A. M. Dirac, *Proc. R. Soc. London*, **113A**, 621 (1927).
6. P. Jordan, *Z. Phys.* **38**, 513 (1926).

7. J. Schwinger, *Quantum Electrodynamics* (Dover Publications, Inc. New York, 1958).
8. R. A Leacock and M. J. Padgett *Phys. Rev. Lett.* **50**, 3 (1983).
9. R. A. Leacock and M. J. Padgett *Phys. Rev.* **D28**, 2491 (1983).
10. R. Courant and D. Hilbert, *Methods of Mathematical Physics*, Vol 1 (Wiley interscience, New York, 1953).
11. E. L. Ince, *Ordinary Differential Equations* (Dover Publications, Inc, New York, 1956).
12. R. S. Bhalla, A. K. Kapoor and P. K. Panigrahi, *Am. J. Phys.* **65**, 1187 (1997).
13. R. S. Bhalla, A. K. Kapoor and P. K. Panigrahi *Mod. Phys. Lett. A*, **12**, 295 (1997).
14. R. S. Bhalla, *The quantum Hamilton - Jacobi approach to energy spectra of potential problems*, *Ph. D. thesis* submitted to University of Hyderabad (1996).
15. F. Cooper, A. Khare and U. Sukhatme, *Supersymmetry in Quantum Mechanics* (World Scientific, Singapore, 2001) and references therein.
16. F. Cooper, A. Khare and U. Sukhatme, *Phys. Rep.* **251**, 267 (1995).

17. R. S. Bhalla, A. K. Kapoor and P. K. Panigrahi, *Int. J. Mod. Phys. A*, **12**, No. **10**, 1875 (1997).
18. S. Jayanthi, *Study of one dimensional potential problems using quantum Hamilton - Jacobi formalism*, M. Phil thesis submitted to University of Hyderabad (1998).
19. A. G. Ushveridze, *Quasi-exactly solvable models in quantum mechanics*, (Institute of Physics Publishing, Bristol, 1994).
20. K. G. Geojo, S. Sree Ranjani and A. K. Kapoor, *J. Phys A : Math. Gen.* **36**, 4591 (2003); preprint quant - ph/0207036.
21. K. G. Geojo, *Quantum Hamilton - Jacobi study of wave functions and energy spectrum of solvable and quasi - exactly solvable models*, Ph. D. thesis submitted to University of Hyderabad (2004).
22. S. Sree Ranjani, K. G. Geojo, A. K Kapoor and P. K. Panigrahi, to be published in *Mod. Phys. Lett. A*. **19**, No. **19**, 1457 (2004); preprint quant - ph/0211168.
23. W. Magnus and S. Winkler, *Hills Equation* (Interscience Publishers, New York, 1966).
24. F. M. Arscot, *Periodic Differential equations* (Pergamon, Oxford, 1964).
25. A. Khare and U. Sukhatme, *J. Math. Phys.* **40**, 5473 (1999); preprint quant - ph/9906044.

- 26. A. Khare and U. Sukhatme, *J. Math. Phys.* **42**, 5652 (2001); preprint quant - ph/0105044.
- 27. A. Khare and B. P. Mandal; quant-ph/0004019.
- 28. B. Bagchi, S. Mallik, C. Quesne and R. Roychoudhury; quant-ph/0107095.
- 29. Z. Ahmed, *Phys. Lett. A.* **282**, 343 (2001).
- 30. F. L. Scarf, *Phys. Rev.* **D57**, 1271 (1998).
- 31. S. Sree Ranjani, A. K. Kapoor and P. K. Panigrahi, to be published in *Mod. Phys. Lett. A.* **19**. No. **27** 2047 (2004); preprint quant - ph/0312041
- 32. S. Sree Ranjani, A. K. Kapoor and P. K. Panigrahi; preprint quant - ph/0403196.
- 33. S. Sree Ranjani, A. K. Kapoor and P. K. Panigrahi; preprint quant - ph/0403054.

Chapter 2

Quantum Hamilton - Jacobi Formalism

In this chapter, we will briefly discuss the classical Hamilton - Jacobi (HJ) formalism, followed by a detailed description of the QHJ theory. By means of an example, the singularity structure of the QMF is explained and we will show, how one can obtain the energy eigenvalues from the exact quantization condition and simultaneously obtain the eigenfunctions. The example, which we use to elucidate the method, is the well known hydrogen atom, for which, we obtain both the eigenvalues and eigenfunctions.

2.1 Classical Hamilton - Jacobi theory

The classical HJ theory deals with continuous canonical transformations, which relates the old canonical coordinates (q_i, p_i) at time t to the new canonical coordinates (Q_i, P_i) , which are constant and may have initial values (q_0, p_0) at time t_0 [1, 2]. The equations of transformations are of the following form

$$q_i = q_i(Q_i, P_i, t_0) \ ; \ p_i = p_i(Q_i, P_i, t_0). \quad (2.1)$$

One can go from the old coordinates to the new coordinates with the help of a generating function. There are four types of generating functions and in our discussion, we consider a generating function $F_2(q_i, P_i, t)$, with q_i and P_i as the independent variables, which will be useful for our present discussion. The new Hamiltonian K is related to the old Hamiltonian H by

$$K = H + \frac{\partial}{\partial t} F_2(q_i, P_i, t). \quad (2.2)$$

If K is equal to zero, then

$$\begin{aligned} \dot{Q}_i &= \frac{\partial K}{\partial P_i} = 0, \\ \dot{P}_i &= -\frac{\partial K}{\partial Q_i} = 0, \end{aligned} \quad (2.3)$$

such that, the new canonical coordinates are constants and (2.2) becomes

$$H + \frac{\partial}{\partial t} F_2(q_i, P_i, t) = 0. \quad (2.4)$$

The transformation equations in terms of the generating function are

$$\begin{aligned} p_i &= \frac{\partial}{\partial q_i} F_2(q_i, P_i, t) \\ Q_i &= \frac{\partial}{\partial P_i} F_2(q_i, P_i, t). \end{aligned} \quad (2.5)$$

Using the above equations in (2.4), one obtains

$$H(q_i, \frac{\partial}{\partial q_i} F_2(q_i, P_i, t), t) + \frac{\partial}{\partial t} F_2(q_i, P_i, t) = 0, \quad (2.6)$$

which is known as the HJ equation, which constitutes a partial differential equation of $F_2(q_i, P_i, t)$ in $(n+1)$ variables. The solution of (2.6) is termed as

the Hamilton's principle function and depends on n constants of integration α_i so that, we may write

$$F_2 \equiv S(q_i, \alpha_i, t) \quad (2.7)$$

and the new momenta P_i can be chosen to be the independent constants of integration α_i , i. e. $P_i = \alpha_i$. Now the transformations in (2.5) can be written as

$$p_i = \frac{\partial}{\partial q_i} S(q_i, \alpha_i, t), \quad (2.8)$$

which is the classical momentum function and

$$Q_i = \frac{\partial}{\partial \alpha_i} S(q_i, \alpha_i, t) = \beta_i. \quad (2.9)$$

where β_i is a constant. At time t_0 , these constitute $2n$ equations relating $n \alpha_i$'s and $n \beta_i$'s to the initial q_i, p_i values, using which, one can evaluate the constants of integration in terms of the initial coordinates. Thus, the above equation can be used to obtain q_i, p_i in terms of α, β and t as

$$q_i = q_i(\alpha_i, \beta_i, t), \quad p_i = p_i(\alpha_i, \beta_i, t). \quad (2.10)$$

Therefore, when solving the HJ equation, one not only obtains the generating function for the canonical transformation but also obtains the solution of the mechanical problem.

When H is not an explicit function of t , then the HJ equation for S can be written as

$$H\left(q_i, \frac{\partial S}{\partial q_i}\right) + \frac{\partial S}{\partial t} = 0, \quad (2.11)$$

where the first term involves only the q dependence and the second term involves time. Hence, one can assume a solution of the form

$$S(q_i, \alpha_i, t) = W(q_i, \alpha_i) - \alpha_1 t, \quad (2.12)$$

where W is the Hamilton's characteristic function. Substituting this in (2.11) gives the HJ equation for the characteristic function as

$$H\left(q_i, \frac{\partial W}{\partial q_i}\right) = \alpha_1. \quad (2.13)$$

The transformation equations will be

$$\begin{aligned} p_i &= \frac{\partial W}{\partial q_i}, \\ Q_i &= \frac{\partial W}{\partial P_i} = \frac{\partial W}{\partial \alpha_i}. \end{aligned} \quad (2.14)$$

Since, H and W are independent of t , from (2.2), one obtains

$$K = \alpha_1 \quad (2.15)$$

and the equations of motion are

$$\dot{P}_i = -\frac{\partial K}{\partial Q_i} = 0 \quad (2.16)$$

with solution $P_i = \alpha_i$ and

$$\begin{aligned} \dot{Q}_i &= \frac{\partial K}{\partial \alpha_1} = 1 \quad i = 1 \\ &= 0 \quad i \neq 1 \end{aligned} \quad (2.17)$$

with solutions

$$\begin{aligned} Q_1 &= t + \beta_1 \equiv \frac{\partial W}{\partial \alpha_1}, \quad i = 1 \\ Q_i &= \beta_i \equiv \frac{\partial W}{\partial \alpha_i}, \quad i \neq 1 \end{aligned} \quad (2.18)$$

From the above equation, we see that only Q_1 is not a constant, but is a function of time. Thus, with HJ equation of the characteristic function, one can solve the systems in which H is independent of t .

2.1.1 Action - angle variables

As a variation to the above discussed HJ method, one can deal with systems with periodic motion. Here, for each periodic motion, one introduces the action variable defined in terms of the momentum p as

$$J = \oint p dq, \quad (2.19)$$

where the integration is carried over a complete period of the periodic motion in the phase space. The Hamilton's characteristic function in terms of this new variable is

$$W = W(q, J). \quad (2.20)$$

The coordinate conjugate of J , known as the angle variable is defined as

$$\omega = \frac{\partial W}{\partial J} \quad (2.21)$$

and the corresponding equation of motion is

$$\dot{\omega} = \frac{\partial H(J)}{\partial J} \equiv \nu \quad (2.22)$$

with the solution

$$\omega = \nu t + \beta \quad (2.23)$$

where ν is the frequency. Equation (2.21) can be solved for q as a function of ω and J , which, in combination with (2.23), will give the desired solution as a function of time.

In order to see how these variables are advantageous, consider the change in ω , as q goes through a complete cycle of rotation, which is given by

$$\Delta\omega = \oint \frac{\partial\omega}{\partial q} dq. \quad (2.24)$$

using (2.21), one gets

$$\Delta\omega = \oint \frac{\partial^2 W}{\partial q \partial J} dq. \quad (2.25)$$

Since J is a constant, one can write

$$\Delta\omega = \frac{d}{dJ} \oint \frac{\partial W}{\partial q} dq \equiv \frac{d}{dJ} \oint p dq = 1, \quad (2.26)$$

where (2.19) has been put to use. From (2.23), it follows that, if τ is the period for complete cycle of q , then

$$\Delta\omega = 1 = \nu\tau. \quad (2.27)$$

Hence, the constant ν can be identified as the reciprocal of the time period

$$\nu = \frac{1}{\tau} \quad (2.28)$$

and therefore, the frequency is associated with the periodic motion of q . Thus for a one dimensional periodic system, the frequency of the system can be found by determining H as a function of J and using (2.22). Hence the use of action - angle variables, provides a powerful technique for obtaining the frequency of the periodic motion, without finding a complete solution of the motion of the system.

2.2 Quantum Hamilton - Jacobi Formalism

The QHJ formalism draws its inspiration from the classical HJ theory and was first initiated in this form by Leacock and Padgett, in 1983 [3,4]. As mentioned in the introduction, the central entity in this formalism is the quantum momentum function $\vec{p}$, which is defined in three dimensions as

$$\vec{p} = \vec{\nabla} S \quad (2.29)$$

where S is analogous to the classical characteristic function W . S is related to the solution of the Schrödinger equation,

$$-\frac{\hbar^2}{2m} \nabla^2 \psi(x, y, z) + V(x, y, z) \psi(x, y, z) = E \psi(x, y, z) \quad (2.30)$$

by the relation

$$\psi(x, y, z) = \exp\left(\frac{iS}{\hbar}\right) \quad (2.31)$$

which, when substituted in (2.30), gives

$$(\vec{\nabla} S)^2 - i\hbar \vec{\nabla} \cdot (\vec{\nabla} S) = 2m(E - V(x, y, z)). \quad (2.32)$$

Substituting (2.29) in (2.32) gives the QHJ equation for $\vec{p}$ as

$$(\vec{p})^2 - i\hbar \vec{\nabla} \cdot \vec{p} = 2m(E - V(x, y, z)) \quad (2.33)$$

and from (2.29) and (2.31), one can see that $\vec{p}$ is the logarithmic derivative of $\psi(x, y, z)$ i. e.

$$\vec{p} = -i\hbar \frac{\vec{\nabla} \psi(x, y, z)}{\psi(x, y, z)} \quad (2.34)$$

The above discussion of the QHJ formalism is done in three dimensions but, from here on, we proceed to one dimension, as we will be discussing only one dimensional potentials in this thesis. One can see that in one dimension, the QHJ equation (2.33) takes the form of the Riccati equation

$$p^2 - i\hbar \frac{dp}{dx} = 2m(E - V(x)). \quad (2.35)$$

In the QHJ formalism, x is treated as a complex variable, there by extending the definition of p to the complex plane. To obtain the full solution of the Riccati equation, one needs to know p completely in the complex plane. From (2.33), one observes that

$$\lim_{\hbar \rightarrow 0} p \rightarrow p_c, \quad (2.36)$$

where $p_c = \sqrt{2m(E - V(x))}$ is the classical momentum, which can be used as a boundary condition on p . When x is a complex variable, p_c needs to be defined carefully, since, it is a double valued function. In the phase space, p_c vanishes at points where $E = V(x)$. In the complex x plane, the classical region is on the real axis, in between the turning points x_1 and x_2 . Correspondingly, p_c is given a branch cut in between the turning points and is defined to be the branch, which is positive just below the branch cut [3, 4]. This completes the definition of p_c and the role of (2.36) is not only that of a boundary condition on p , but also as a correspondence principle.

2.2.1 Singularities of the QMF

It has already been mentioned that the singularity structure of the QMF plays an important role in obtaining the eigenfunctions and eigenvalues through

the QHJ formalism. The location of singularities of p and the residues at these singular points are the essential ingredients required in this formalism.

The QMF has two types of singularities, the moving and the fixed singularities. It is well known from the oscillation theorem, that the n^{th} excited state wave function has n zeros on the real line. These zeros correspond to the poles of p , as seen from (2.34). The location of these poles depends on the energy and the initial conditions. Hence, these are called as moving poles. For Riccati equation, it is known that only poles can appear as moving singularities [5]. It is easy to see that the residue at these moving poles is $-i\hbar$. Let x_0 be a point, where $V(x)$ is analytic and p has a moving pole. Assuming a moving pole of order m , we proceed to calculate the residue of p at x_0 , by expanding p in Laurent expansion around x_0 as

$$p = \sum_{k=1}^m \lambda_k (x - x_0)^{-k} + \sum_{k=0}^{\infty} \gamma_k (x - x_0)^k. \quad (2.37)$$


Substituting (2.37) in the QHJ equation (2.35) and comparing the coefficients of different powers of $x - x_0$, one gets $m = 1$ and

$$\lambda_1 = -i\hbar. \quad (2.38)$$

Thus, the moving poles are simple poles with residue $-i\hbar$. This information of the moving poles is used to obtain the quantization condition satisfied by the quantum action

$$J = J(E) \equiv \frac{1}{2\pi} \oint_C p dx. \quad (2.39)$$

Here, C is the contour, which encloses the moving poles of p as shown in the fig. 2. 1.

**Fig. 2. 1**

Applying the Cauchy residue theorem to the n poles enclosed by C , one obtains the quantization condition

$$J(E) \equiv \frac{1}{2\pi} \oint_C p dx = n\hbar, \quad (2.40)$$

which can be viewed as a generalization of the Wilson - Sommerfeld quantization. Unlike the WKB quantization and the Wilson - Sommerfeld quantization, this is an exact quantization condition and is a result of the oscillation theorem and is true only for the bound states [5, 6]. Hence one cannot apply (2.40) to continuous energy states with wave functions non vanishing at infinity.

The power of the classical HJ theory lies in the fact that one obtains the frequencies of the system directly from the classical action - angle variables.

Analogously, in the QHJ formalism, one can obtain the energy eigenvalues of the stationary states directly from the quantum action variable making use of the exact quantization condition (2.40). To arrive at the energy eigenvalues, one needs to know the location of other singularities of the QMF, which lie outside the contour C and their corresponding residues. The other singularities of the QMF arise from the potential $V(x)$ and are called as fixed singularities. The locations of these singular points are energy independent and appear in all the solutions of the QHJ equation. Thus, if the potential is meromorphic, the QMF will also be meromorphic. The residue at any of these fixed poles can also be computed by using the QHJ equation (2.33). Due to the quadratic nature of the QHJ equation, one obtains two values for each of these residues at the fixed poles. Of these two values, one needs to check which of the values satisfy the boundary condition (2.36). Since, p_c is defined to be positive just below the branch cut, we choose those values of residues, which in the limit $\hbar \rightarrow 0$, correspond to positive p_c below the branch cut. This method of choosing the residue is explained in the study of hydrogen atom, later in the chapter.

In all our investigations, we have assumed that the QMF has only finite number of singularities in the complex plane, which turned out to be true for all the ES and QES models studied. This assumption is equivalent to saying that the point at infinity is an isolated singularity. This property is shared by both the ES and QES models and the only difference is the location of the moving poles. For ES models, the QMF has poles only on the real line,

whereas for QES models, the moving poles are in both real and complex locations. In case of periodic potentials, it will be shown that both ES and QES cases share the same kind of singularity structure. The knowledge of the singularity structure allows one to write the QMF as a function of x and hence compute the eigenvalues and the eigenfunctions. The integral $J(E)$ in (2.40) can be computed in terms of the residues at the singularities, outside the contour C . This gives the energy eigenvalues for one dimensional ES models [7] and for QES models, one is lead to the QES [8, 9] condition by the same condition.

In the next section, we clearly explain all the above discussed points using the example of the hydrogen atom. We also show directly from the QHJ formalism that without using the quantization condition, we can arrive at the form of the QMF and hence, the values of the eigenvalues and eigenfunctions by using well known results in complex variables.

2.3 Hydrogen atom

In this section, we obtain the bound state wave functions for the radial part of the Coulomb problem [10]. We consider the supersymmetric potential

$$V_-(r) = -\frac{e^2}{r} + \frac{l(l+1)}{r^2} + \frac{e^4}{4(l+1)^2} \quad (2.41)$$

which is equal to the original coulomb potential plus the constant $\frac{e^4}{4(l+1)^2}$, which has been added to make the ground state energy equal to zero. The

radial Schrödinger equation with $\hbar = 2m = 1$ is given by

$$\frac{d^2 R}{dr^2} + \frac{2}{r} \frac{dR}{dr} + \left(E + \frac{e^2}{r} - \frac{l(l+1)}{r^2} - \frac{e^4}{4(l+1)^2} \right) R = 0. \quad (2.42)$$

Using the transformation $R(r) = u(r)/r$, $p = -i(u'(r)/u(r))$, one obtains the QHJ equation as

$$p^2 - i \frac{dp}{dr} - \left(E + \frac{e^2}{r} - \frac{l(l+1)}{r^2} - \frac{e^4}{4(l+1)^2} \right) = 0. \quad (2.43)$$

The wave function $u(r)$ should vanish at $r = 0$ and hence the QMF has fixed pole at the origin along with the n moving poles with residue $-i$ on the positive real line. We assume that there are no other singular points of p in the complex plane, which makes *the point at infinity, an isolated singularity of p* . With this singularity structure of p , we, now proceed to find the energy eigenvalues.

From this information about the singularity structure of the QMF, we know that p is meromorphic and has the above mentioned singular points. Hence, one can write p as a sum of singular and analytic parts as written below

$$p = \sum_{k=1}^n \frac{-i}{r - r_k} + \frac{b_1}{r} + Q(r). \quad (2.44)$$

In the above equation, the summation term describes the sum of all the principal parts coming from all the individual Laurent expansions of p taken around each moving pole with $-i$ as the residue. The second term in (2.44) represents the singular part in the Laurent expansion of p around $r = 0$, with b_1 as the residue and $Q(r)$ represents the analytic part of p . From the QHJ

equation (2.43), one can see that for large r , p goes to zero. Thus $Q(r)$ is an entire function and bounded at infinity. Thus, from Louville's theorem, $Q(r)$ is a constant, say C . Thus, (2.44) becomes

$$p = \sum_{k=1}^n \frac{-i}{r - r_k} + \frac{b_1}{r} + C, \quad (2.45)$$

which gives the form of p in the entire complex plane.

To obtain the energy eigenvalues one only needs to calculate the value of the residue b_1 and the value of the constant C . On the other hand, the location r_k of the moving poles must be found in order to obtain the wave function.

2.3.1 Definition of p_c

As mentioned in the previous section, due to the quadratic nature of the QHJ equation, b_1 will have two values. To select the right value of residue, we make use of the boundary condition (2.36), which uses the classical momentum p_c . Hence, we first proceed to define p_c which is given by

$$p_c = \sqrt{E + \frac{e^2}{r} - \frac{l(l+1)}{r^2} - \frac{e^4}{4(l+1)^2}} \quad (2.46)$$

in the complex plane and which is equivalent to

$$p_c \equiv \frac{\pm i\alpha}{2r} \left(r^2 + \frac{e^2 r}{\alpha^2} - \frac{l(l+1)}{\alpha^2} \right)^{1/2}. \quad (2.47)$$

where $\alpha = \sqrt{\frac{e^4}{4(l+1)^2} - E}$, is a positive constant. Let r_1 and r_2 be the two turning points and in the complex r plane, p_c is given a branch cut between the two turning points. p_c is defined to be positive just below the branch cut.

Hence, one needs to choose that branch of p_c from (2.47) which satisfies this condition. In order to be able to do that we introduce the variables, ρ_1, ρ_2, θ_1 and θ_2 such that

$$r - r_1 = \rho_1 \exp(i\theta_1), \quad r - r_2 = \rho_2 \exp(i\theta_2) \quad (2.48)$$

where θ_1 and θ_2 are the angles as shown in the fig 2.2 below




Fig. 2. 2

and are chosen to lie in the open interval $(0, 2\pi)$. This gives a branch cut to p_c , for $r_1 < r < r_2$. Thus, one can write p_c as

$$p_c \sim \frac{i\lambda\alpha}{2r} \sqrt{(r - r_1)(r - r_2)}, \quad (2.49)$$

where $\lambda = \pm 1$ and using (2.48), p_c becomes

$$p_c \equiv \frac{i\lambda\alpha}{2r}(\rho_1\rho_2)^{1/2} \exp\left(i\frac{(\theta_1 + \theta_2)}{2}\right). \quad (2.50)$$

We need to fix the sign of λ , which will fix the branch of p_c to be selected. For this, we consider a point just below the branch cut which corresponds to $\theta_1 \sim 2\pi$, $\theta_2 \sim \pi$, which gives

$$p_c = \frac{\lambda\alpha}{2r}(\rho_1\rho_2)^{1/2}. \quad (2.51)$$

Therefore, for p_c to be positive just below the branch cut, we choose $\lambda = +1$. Thus, we obtain the definition of p_c in the entire complex plane as

$$p_c = \frac{i\alpha}{2r}(\rho_1\rho_2)^{1/2} \exp\left(i\frac{(\theta_1 + \theta_2)}{2}\right), \quad (2.52)$$

with $0 < \theta_1 < 2\pi$ and $0 < \theta_2 < 2\pi$.

2.3.2 Residue at the fixed pole $r = 0$

With the complete definition of p_c obtained, we proceed to calculate the residue of both p_c and p at $r = 0$. Comparing the residues, one can fix the correct value of b_1 . First, we calculate the residue of p_c . At the origin $\theta_1 \sim \pi$ and $\theta_2 \sim \pi$, which, when substituted in (2.52), give the residue at $r = 0$ as (2.52) p_c at $r = 0$

$$-\frac{i\alpha}{2}(r_1r_2)^{1/2}. \quad (2.53)$$

Now to calculate the residue of p , we perform a Laurent expansion of p around $r = 0$ as

$$p = \frac{b_1}{r} + a_0 + a_1r + \dots \quad (2.54)$$

Substituting (2.54) in (2.43) and comparing the coefficients of $1/r^2$, one obtains the quadratic equation

$$b_1^2 + ib_1 + l(l+1) = 0, \quad (2.55)$$

which gives the two values of b_1 as

$$b_1 = il, -i(l+1). \quad (2.56)$$

Comparing the values of b_1 with the residue of p_c in (2.53), we see that the sign of the imaginary part must be negative. Hence, we must select

$$b_1 = -i(l+1), \quad (2.57)$$

which satisfies the boundary condition (2.36).

2.3.3 Calculation of energy eigenvalues and eigenfunctions

With the selection of the right value of the residue at the fixed pole, one can write p as

$$p = -i \frac{P'_n(r)}{P_n(r)} - \frac{i(l+1)}{r} + C \quad (2.58)$$

where $\sum_{k=1}^n \frac{-i}{r-r_k} = -i \frac{P'_n(r)}{P_n(r)}$ and $P_n(r)$ is an n^{th} degree polynomial. Substituting p from (2.58) in (2.43) one obtains

$$\begin{aligned} \frac{P''_n(r)}{P_n(r)} - 2iC \left(\frac{P'_n(r)}{P_n(r)} + \frac{(l+1)}{r} \right) - \frac{2(l+1)}{r} \frac{P'_n(r)}{P_n(r)} \\ + C^2 - E - \frac{e^2}{r} + \frac{e^4}{4(l+1)^2} = 0. \end{aligned} \quad (2.59)$$

In order to fix C and obtain the energy eigenvalues, we look at the behaviour of each term of (2.59), for large r . Noting that for large r ,

$$\frac{P_n''(r)}{P_n(r)} \sim \frac{n(n-1)}{r^2} ; \quad \frac{P_n'(r)}{P_n(r)} \sim \frac{n}{r}, \quad (2.60)$$

we equate the coefficients of the constant terms and coefficients of $1/r$ in (2.59) to zero individually and get

$$C = \frac{ie^2}{2(n+l+1)}; \quad E = C^2 + \frac{e^4}{4(l+1)^2} \quad (2.61)$$

which in turn give

$$E = -\frac{e^4}{4(n+l+1)^2} + \frac{e^4}{4(l+1)^2}, \quad (2.62)$$

which is the required energy eigenvalue expression and one is left with the second order differential equation

$$P_n''(r) + P_n'(r) \left(\frac{2(l+1)}{r} - \frac{e^2}{n+l+1} \right) + e^2 \left(\frac{1}{r^2} - \frac{l+1}{(n+l+1)r} \right) P_n(r) = 0. \quad (2.63)$$

Defining $\frac{e^2}{n+l+1}r = y$ in (2.63), we get

$$yP_n''(y) + ((2l+1) + 1 - y)P_n'(y) + nP_n(y) = 0 \quad (2.64)$$

which is the associated Laguerre equation, and $P_n(y)$ is proportional to the Laguerre polynomial, denoted by $L_n^{2l+1}(y)$. The bound state wave function can be obtained by using (2.34), which gives

$$u(r) = \exp \left(i \int p dx \right) \quad (2.65)$$

Substituting the value of p from (2.58) in the above equation and changing the variable from r to y , one obtains the wave function in terms of y as

$$u_n(y) = y^{l+1} \exp\left(-\frac{y}{2}\right) L_n^{2l+1}(y), \quad (2.66)$$

which matches with the known result [11].

We would like to mention here that one can obtain the energy eigenvalues by using the exact quantization condition (2.40) discussed in §2.2 without solving the QHJ equation. For this, one needs to calculate the residue at infinity, which will again have two values. The right value is chosen by applying the boundary condition (2.36) and by using the Cauchy residue theorem, one is led to the expression for the energy eigenvalues, the details of which are given in [12]. Instead of using the above mentioned boundary condition, one may note that p (2.45) is a rational function of x and hence, the sum of all the residues (including the residue at infinity) is zero. This is equivalent to what one gets from the exact quantization condition and is widely used in our analysis of periodic and PT symmetric potentials.

2.4 Change of variables in QHJ Formalism

We would like to emphasize here that, locating the singular points of the QMF and imposing the boundary condition (2.36) are two crucial steps in arriving at the correct physical solutions. For many problems, it is necessary to perform a change of variable

$$y = f(x). \quad (2.67)$$

Thus, for a general potential $V(x)$, the QHJ equation (2.35) with $\hbar = 2m = 1$ is

$$p^2 - i \frac{dp}{dx} - (E - V(x)) = 0, \quad (2.68)$$

which, after a change of variable (2.67) and introduction of q by $q(y) \equiv ip(x(y))$, becomes

$$q^2(y) + F(y) \frac{dq(y)}{dy} + E - \tilde{V}(y) = 0 \quad (2.69)$$

Here $F(y)$ equals the derivative $\frac{df(x)}{dx}$ expressed as a function of y and $\tilde{V}(y) = V(x(y))$. In order to calculate the residue at the moving poles, removing the complexity caused due to the presence of $F(y)$, we bring (2.69) to a simple form, by introducing χ through the following transformation equations.

$$q = F(y)\phi, \quad \chi = \phi + \frac{1}{2} \frac{F'(y)}{F(y)}, \quad (2.70)$$

when substituted in (2.69), give

$$\chi^2 + \frac{d\chi}{dy} + \frac{E - \tilde{V}(y)}{(F(y))^2} - \frac{1}{2} \left(\frac{F''(y)}{F(y)} \right) + \frac{1}{4} \left(\frac{F'(y)}{F(y)} \right)^2 = 0, \quad (2.71)$$

where, the residue at moving poles is one. The function χ will also be called as the QMF and equation (2.71) as the QHJ equation. From (2.71), one can again see that the residue at the moving poles of χ is unity. The expression for the wave function, in the new variable y can be obtained as follows. From (2.34), one obtains ψ with $\hbar = 1$ as

$$\psi(x) = \exp \left(i \int p dx \right) \quad (2.72)$$

in which, substituting $p = -iq$, followed by the change of variable (2.67) gives

$$\psi(y) = \exp \left(\int \frac{q}{F(y)} dy \right). \quad (2.73)$$

Using the transformation equations given in (2.70), one is led to the expression for $\psi(y)$ in terms of χ and $F(y)$ as

$$\psi(y) = \exp \int \left(\chi - \frac{1}{2} \frac{d}{dy} (\ln F(y)) \right) dy. \quad (2.74)$$

It is not always easy to impose the boundary condition (2.36). We, therefore, describe a set of constraints, which will be found useful in restricting the choice of residues at the fixed poles. Firstly, the wave functions should be finite every where for the physical values of x . If one is looking for bound state solutions, one must have the condition that the wave function vanishes for large x . Since, in this thesis, we are dealing only with the one dimensional potentials, we can also use the fact that vanishing of wave function for large x , implies that the bound state levels are non - degenerate. So, for example, parity invariance will imply $\psi(-x) = \pm\psi(x)$ and QMF will satisfy $p(-x) = -p(x)$. Also the bound state wave functions will be real and hence $p^*(x) = p(x)$. All possible values of the residues, consistent with these and other similar conditions, must be accepted. In all the cases studied here, the QMF turns out to be a rational function and hence, the quantization condition becomes equivalent to *the sum of all the residues of QMF, including the residue at infinity, being equal to zero*. This leads to a relation between the residues and the number of moving poles n . As we will see in later chapters,

some combinations of residues get eliminated because the number n has to be positive.

References

1. H. Goldstein, *Classical Mechanics* (Addison-Wesley/Narosa, Indian student edition, 1995)
2. E. J. Saletan and A. H. Cromer, *Theoretical Mechanics* (John Wiley, New York, 1971).
3. R. A Leacock and M. J. Padgett, *Phys. Rev. Lett.* **50**, 3 (1983).
4. R. A. Leacock and M. J. Padgett, *Phys. Rev.* **D28**, 2491 (1983).
5. E. L. Ince, *Ordinary Differential Equations* (Dover Publications, Inc, New York, 1956).
6. R. Courant and D. Hilbert, *Methods of Mathematical Physics*, Vol 1 (Wiley interscience, New York, 1953).
7. S. Sree Ranjani, K. G. Geojo, A. K Kapoor and P. K. Panigrahi, to be published in *Mod. Phys. Lett. A*, **19**, No. **19**, 1457 (2004); preprint quant - ph/0211168.
8. K. G. Geojo, S. Sree Ranjani and A. K. Kapoor, *J. Phys. A : Math. Gen.* **36**, 4591 (2003); quant - ph/0207036.
9. K. G. Geojo, *Quantum Hamilton - Jacobi study of wave functions and energy spectrum of solvable and quasi - exactly solvable models*, Ph. D. thesis submitted to University of Hyderabad (2004).

10. L. I. Schiff, *Quantum Mechanics* (McGraw - Hill, Singapore, 1986).
11. F. Cooper, A. Khare and U. Sukhatme, *Supersymmetry in Quantum Mechanics* (World Scientific, Singapore, 2001) and references therein.
12. R. S. Bhalla, *The quantum Hamilton - Jacobi approach to energy spectra of potential problems*, *Ph. D. thesis* submitted to University of Hyderabad (1996).

Chapter 3

Study of potentials exhibiting different spectra

3.1 Introduction

This chapter is devoted to the study of potentials, which exhibit different spectra for different ranges of the potential parameters. The first potential investigated, here, is the Scarf - I potential, which has been well studied [1, 2] and is known to exhibit both phases of SUSY, namely the broken and the unbroken phases of SUSY, for different parameter ranges. In [3], Bhalla *et. al.* have studied this potential using the QHJ formalism and obtained the the energy eigenvalue expressions in both the phases. In the present study, we obtain both the eigenvalues and eigenfunctions along with the different ranges.

The second potential to be discussed is the Scarf potential[4, 5], which exhibits band structure in one range and ordinary bound state spectrum in the other range. In [5], a group theoretical treatment of Scarf potential has been discussed with the bound state and scattering state problem, being

extended to include the band structure. Using the QHJ formalism, we show that one can arrive at the bound state solutions and the band edge solutions of this potential in a simple and straight forward fashion. The next section of the chapter discusses the Scarf - I potential, followed by a section on Scarf potential.

3.2 Phases of Supersymmetry

The expression for the supersymmetric Scarf - I potential is

$$V_-(x) = -A^2 + (A^2 + B^2 - A\alpha\hbar) \sec^2 \alpha x - B(2A - \alpha\hbar) \tan \alpha x \sec \alpha x \quad (3.1)$$

and it exhibits broken and unbroken phases of SUSY for the following ranges of potential parameters [1]. In the parameter range,

$$(A - B) > 0, (A + B) > 0, \quad (3.2)$$

SUSY is exact and SUSY is broken for the range

$$(A - B) > 0, (A + B) < 0. \quad (3.3)$$

The supersymmetric phases of this potential are well studied and their corresponding bound state solutions are available in the literature [1-3]. In this chapter, we will show that the ranges of the two different phases and their corresponding solutions emerge naturally from the QHJ analysis [6].

Putting $2m = 1$ and writing the QHJ equation in terms of $q = ip$ we get

$$q^2 + \hbar q' + E + A^2 - (A^2 + B^2 - A\alpha\hbar) \sec^2 \alpha x + B(2A - \alpha\hbar) \tan \alpha x \sec \alpha x = 0 \quad (3.4)$$

As discussed in the end of §2.4, we perform a change of variable

$$y = \sin \alpha x, \quad (3.5)$$

which gives $F(y) = \alpha\sqrt{1-y^2}$. The transformation equations (2.70) of §2.4 take the form,

$$q = \alpha\sqrt{1-y^2}\phi ; \quad \phi = \chi + \frac{y\hbar}{2(1-y^2)}. \quad (3.6)$$

and the QHJ equation in terms of χ is obtained as,

$$\begin{aligned} \chi^2 + \hbar\chi' + \frac{y^2\hbar^2}{4(1-y^2)^2} + \frac{E + A^2}{\alpha^2(1-y^2)} &+ \frac{\alpha^2\hbar - 2(A^2 + B^2 - A\alpha\hbar)}{2\alpha^2(1-y^2)^2} \\ &+ \frac{B(2A - \alpha\hbar)y}{\alpha^2(1-y^2)^2} = 0. \end{aligned} \quad (3.7)$$

3.2.1 Form of QMF (χ)

From (3.7), one can see that χ has fixed poles at $y = \pm 1$. In addition, there will be n moving poles, with residue one, on the real line corresponding to the n nodes of the n^{th} excited state. Proceeding in the same way as in the case of hydrogen atom, we assume that χ has only the above mentioned singularities in the complex plane. Hence, we write χ as

$$\chi = \frac{b_1}{y-1} + \frac{b'_1}{y+1} + \hbar \frac{P'_n(y)}{P_n(y)} + C, \quad (3.8)$$

where b_1 and b'_1 are the residues at $y = \pm 1$ respectively and $P_n(y)$ is an n^{th} degree polynomial. C represents the analytic part of χ which is a constant due to Liouville's theorem, since χ is bounded at infinity as seen from (3.7). As in §2.3, we first define the classical momentum function, which will help us to fix the right form of χ .

3.2.2 Definition of the Classical momentum function

The classical momentum function, which is given a branch cut in between the two turning points x_1 and x_2 in the complex x plane, is given by

$$p_c = \sqrt{2m(E - V(x))} \quad (3.9)$$

and is defined to be the branch, which is positive just below the branch cut. Since all the analysis is performed in the complex y plane with χ as the QMF, which has fixed poles at $y = \pm 1$, we use the condition

$$\lim_{\hbar \rightarrow 0} \chi \rightarrow \chi_c \quad (3.10)$$

instead of (2.36). The expression of χ_c is obtained as,

$$\chi_c = \frac{i\tilde{p}_c}{\alpha\sqrt{1-y^2}} - \frac{y\hbar}{2(1-y^2)}, \quad (3.11)$$

by putting $q = ip$ in the transformations (3.6) and replacing p with $\tilde{p}_c$, which represents the classical momentum function (3.20) in terms of y as

$$\tilde{p}_c = \pm i \frac{\sqrt{E + A^2}}{\sqrt{1-y^2}} \left(y^2 + \frac{B^2 - A\alpha - E - B(2A - A\alpha\hbar)y}{E + A^2} \right)^{1/2}, \quad (3.12)$$

where $1 - y^2$ is positive as y lies between ± 1 . Let y_1 and y_2 be the images of the two turning points x_1 and x_2 in the y plane. The mapping from $y = \sin \alpha x$ in (3.5), is such that it maps the point just below the branch cut in the x plane to a point just below the branch cut in the y plane. Hence, we select the branch of $\tilde{p}_c$, which is positive just below the branch cut. Similar to the hydrogen atom case, we introduce the variables θ_1, r_1 and θ_2, r_2 , where

$$y - y_1 = r_1 \exp i\theta_1 \quad y - y_2 = r_2 \exp i\theta_2 \quad (3.13)$$

as represented in the fig.3.1.




Fig. 3.1

The values of θ_1 and θ_2 are chosen to lie in the interval $(0, 2\pi)$ and $\tilde{p}_c$ can be written in terms of the above variables as

$$\tilde{p}_c = i\lambda \frac{\sqrt{E + A^2}}{\sqrt{1 - y^2}} (r_1 r_2)^{1/2} \exp \left(i \frac{(\theta_1 + \theta_2)}{2} \right), \quad (3.14)$$

where $\lambda = \pm 1$. Proceeding in the same way as done for the Coulomb problem, we see that for a point just below the branch cut, $\theta_1 \sim 2\pi$ and $\theta_2 \sim \pi$. Substituting these values in (3.14), one gets

$$\tilde{p}_c \sim \frac{\lambda}{\sqrt{1 - y^2}} (E + A^2)^{1/2} (r_1 r_2)^{1/2}. \quad (3.15)$$

For $\tilde{p}_c$ to be positive just below the branch cut, we need to fix $\lambda = 1$. Thus in the entire complex y plane,

$$\tilde{p}_c = i \frac{\sqrt{E + A^2}}{\sqrt{1 - y^2}} (r_1 r_2)^{1/2} \exp \left(i \frac{(\theta_1 + \theta_2)}{2} \right), \quad (3.16)$$

which in turn gives

$$\chi_c = - \frac{\sqrt{E + A^2}}{1 - y^2} (r_1 r_2)^{1/2} \exp \left(i \frac{(\theta_1 + \theta_2)}{2} \right) - \frac{y\hbar}{2(1 - y^2)}. \quad (3.17)$$

3.2.3 Residues at the fixed poles $y = \pm 1$

At $y = -1$: We calculate the residue of χ_c at the fixed pole $y = -1$. In the complex plane the point $y = -1$ corresponds to $\theta_1 \sim \theta_2 \sim \pi$, which gives

$$\chi_c = \frac{\sqrt{E + A^2}}{1 - y^2} (r_1 r_2)^{1/2} - \frac{y\hbar}{2(1 - y^2)}. \quad (3.18)$$

The residue of χ_c at $y = -1$ is

$$\sim \frac{\sqrt{E + A^2}}{2} (r_1 r_2)^{1/2} + \frac{\hbar}{4}, \quad (3.19)$$

which is a positive quantity. Now, we evaluate the residue of χ at $y = -1$, by doing Laurent expansion of χ around $y = -1$, as

$$\chi = \frac{b'_1}{y + 1} + a_0 + a_1(y + 1) + \dots \quad (3.20)$$

Substituting this in (3.7) and comparing the coefficients of $\frac{1}{(y+1)^2}$, one obtains the quadratic equation

$$(b'_1)^2 - \hbar b'_1 + \frac{3\alpha^2 \hbar^2 - (2A + 2B)^2 + 4(A + B)\alpha \hbar}{16\alpha^2} = 0, \quad (3.21)$$

which gives the two values of b'_1 as

$$b'_1 = \frac{(A+B)}{2\alpha} + \frac{\hbar}{4}, \quad -\frac{(A+B)}{2\alpha} + \frac{3\hbar}{4}. \quad (3.22)$$

Since the residue of χ_c at $y = -1$ is positive, (3.10), implies that we select the value of b'_1 , which is positive in the limit $\hbar \rightarrow 0$. Thus, we pick

$$b'_1 = \begin{cases} \frac{A+B}{2\alpha} + \frac{\hbar}{4} & \text{if } (A+B) > 0 \\ -\frac{A+B}{2\alpha} + \frac{3\hbar}{4} & \text{if } (A+B) < 0. \end{cases} \quad (3.23)$$

From the above equation, we see that both the values of b'_1 are acceptable, but in two different ranges. Comparison of these ranges with the ranges given in (3.2) and (3.3), show that they correspond to the different phases of SUSY.

At $y = 1$:

Repeating the above process at the other fixed pole, one obtains the residue of χ_c at $y = 1$ as

$$\sim \frac{\sqrt{E+A^2}}{2}(r_1 r_2)^{1/2} + \frac{\hbar}{4}, \quad (3.24)$$

which is positive. The two values of the residue of χ are

$$b_1 = \frac{(A-B)}{2\alpha} + \frac{\hbar}{4}, \quad -\frac{(A-B)}{2\alpha} + \frac{3\hbar}{4}. \quad (3.25)$$

Comparing equations (3.24) and (3.25), one sees that each value of b_1 is acceptable in different ranges, *i.e*

$$b_1 = \begin{cases} \frac{A-B}{2\alpha} + \frac{\hbar}{4} & \text{if } (A-B) > 0 \\ -\frac{A-B}{2\alpha} + \frac{3\hbar}{4} & \text{if } (A-B) < 0 \end{cases} \quad (3.26)$$

But, the present case of interest is $(A - B) > 0$, for which the choice of residue is

$$b_1 = \frac{(A - B)}{2\alpha} + \frac{\hbar}{4}. \quad (3.27)$$

The range $(A - B) > 0$ corresponds to both phases of SUSY, as seen from (3.2) and (3.3). Thus, from the above discussion, one can see that by applying the boundary condition (3.10) properly, one can obtain both the ranges of the parameters and simultaneously select the values of the residues which give physically acceptable solutions. We now proceed to obtain the eigenvalues and eigenfunctions for both the phases.

3.2.4 Case 1 : $(A - B) > 0$, $(A + B) > 0$

For the above range of parameters, χ in (3.8) becomes

$$\chi(y) = \left(\frac{A - B}{2\alpha} + \frac{\hbar}{4} \right) \frac{1}{y - 1} + \left(\frac{A + B}{2\alpha} + \frac{\hbar}{4} \right) \frac{1}{y + 1} + \hbar \frac{P'_n(y)}{P_n(y)} + C. \quad (3.28)$$

where, the suitable values of b'_1 and b_1 are taken from (3.23) and (3.27) respectively. Substituting χ from the above equation in (3.7) and putting $\hbar = 1$, we compare the constant terms and the coefficients of $1/y^2$ terms individually, in the limit of y going to infinity and obtain $C = 0$ and the energy eigenvalue as

$$E_n = (A + n\alpha)^2 - A^2. \quad (3.29)$$

The equation for $P_n(y)$ assumes the form

$$(1 - y^2)P'' + (\mu - \nu - (\mu + \nu + 1)y)P' + n(n + \mu + \nu)P_n(y) = 0 \quad (3.30)$$

with

$$\mu = \frac{A - B}{\alpha}, \quad \nu = \frac{A + B}{\alpha}, \quad (3.31)$$

is seen to be the Jacobi differential equation and hence $P_n(y)$ coincides with $P_n^{\mu-1/2, \nu-1/2}(y)$ and the bound state wave function for the phase in which SUSY is exact, is

$$\psi_n(y) = N(1 - y)^{\mu/2}(1 + y)^{\nu/2}P_n^{\mu-\frac{1}{2}, \nu-\frac{1}{2}}(y). \quad (3.32)$$

The energy eigenvalues in (3.29) and the wave functions agree with the solutions given in [1].

3.2.5 Case 2 : $(A - B) > 0, \quad (A + B) < 0$

The right choice of the residues for this range, as shown in §3. 2. 3 is,

$$b_1 = \frac{(A - B)}{2\alpha} + \frac{\hbar}{4}, \quad b'_1 = -\frac{(A + B)}{2\alpha} + \frac{3\hbar}{4}. \quad (3.33)$$

Substituting χ from (3.8), with the above values of the residues in (3.7) with $\hbar = 1$, one obtains the energy eigenvalue as

$$E = \left(B - \left(n + \frac{1}{2} \right) \alpha \right)^2 - A^2 \quad (3.34)$$

and the differential equation for $P_n(y)$ turns out to be

$$(1 - y^2)P_n''(y) + ((\nu - \mu + 1) - y(\nu + \mu + 2))P_n'(y) + n(n + \mu + \nu + 1)P_n(y) = 0 \quad (3.35)$$

and the bound state wave function is found to be

$$\psi_n(y) = N(1 - y)^{\frac{\mu}{2}}(1 + y)^{\frac{\nu}{2} + \frac{1}{2}}P_n^{\nu + \frac{1}{2}, \mu - \frac{1}{2}}(y) \quad (3.36)$$

where

$$\mu = \frac{A - B}{\alpha}, \quad \nu = \frac{1}{2} - \frac{A + B}{\alpha} \quad (3.37)$$

The Scarf - I potential can be related to the Pöschl - Teller - I potential by redefinition of the potential parameters as

$$x \rightarrow x + \frac{\pi}{2\alpha}, \quad A = \delta + \beta, \quad B = \delta - \beta. \quad (3.38)$$

With this redefinition of the parameters, we see that the bound state wave functions in (3.36) match with the bound state wave functions of the broken SUSY phase of the Pöschl - Teller - I potential given in [7].

Thus, we see that the QHJ formalism in one dimension gives the accurate expressions for the bound state wave functions, when there are different phases of SUSY. It may be remarked here that, in the range $A - B < 0$, $A + B < 0$, SUSY is exact but the roles of H_- and H_+ are interchanged. In the range $A - B > 0$, $A + B < 0$, SUSY is again broken.

3.3 Phases of a periodic potential

In this section, we study the Scarf potential

$$V(x) = -\frac{V_0}{\sin^2\left(\frac{\pi x}{a}\right)} \quad (3.39)$$

with a as the potential period. The Schrödinger equation for this potential, with $\hbar = 1$, is

$$\frac{d^2\psi(x)}{dx^2} + \frac{\pi^2}{a^2} \left(\lambda^2 + \frac{\left(\frac{1}{4} - s^2\right)}{\sin^2\left(\frac{\pi x}{a}\right)} \right) \psi(x) = 0 \quad (3.40)$$

with λ and s , defined by

$$\lambda^2 = \frac{2mEa^2}{\pi^2}; \quad \left(\frac{1}{4} - s^2\right) = \frac{2mV_0a^2}{\pi^2}. \quad (3.41)$$

This potential is known to exhibit two different types of spectra for different ranges of s , unlike the Scarf - I potential, which exhibits different bound state spectra for different ranges. In the range $s > 1/2$, one can see from the plot in fig. 3.2

with $s = 2$ and $a = 1$, allows ordinary bound stat




Fig. 3. 2

that the potential is an array of infinite potential wells. Hence, the particle is confined to one well and the wave function should vanish at $x = \pm a$. Thus, in the above range, the potential has bound state spectrum. From the above plot, one can also see that the minima of the potential is at zero and hence the energy eigenvalues are always greater than zero. For s , in the range $0 < s < 1/2$, one can see from the plot in fig 3.3,




Fig. 3. 3

that the potential is similar to that of the potential in a crystal lattice and hence there are bands of allowed ranges. The particle can escape to infinity and the wave function need not vanish at $x = \pm a$, but it should not become infinity anywhere. Unlike the previous case one cannot make any statement on the energy eigenvalues just by looking at the plot of the potential. But in the following sections, we will see that the energy is greater than zero.

For this potential, we will show that one can obtain both the ranges and their corresponding solutions, by doing a QHJ analysis. The corresponding QHJ equation for the Scarf potential with $p = -iq$ is

$$q^2 + q' + \frac{\pi^2}{a^2} \left(\lambda^2 + \frac{(\frac{1}{4} - s^2)}{\sin^2(\frac{\pi x}{a})} \right) = 0. \quad (3.42)$$

Performing the change of variable

$$y = \cot \left(\frac{\pi x}{a} \right), \quad (3.43)$$

we make use of the results in §2.4. The expression for df/dx , expressed in y , becomes $F(y) = -\pi(1 + y^2)/a$, which when substituted in (2.70) gives the

transformation equations,

$$q = -\frac{\pi}{a}(1+y^2)\phi ; \quad \phi = \chi - \frac{y}{1+y^2}. \quad (3.44)$$

Thus, the QHJ equation (2.71) for χ , becomes

$$\chi^2 + \frac{d\chi}{dy} + \frac{\lambda^2 - 1}{(y^2 + 1)^2} + \frac{(\frac{1}{4} - s^2)}{y^2 + 1} = 0. \quad (3.45)$$

3.3.1 Form of the QMF χ

The QMF has n moving poles with residue one on the real line, corresponding to the nodes of the n^{th} excited state wave function. From (3.45), one can see that χ has fixed poles at $y \pm i$. Hence, one can write χ in the rational form as

$$\chi = \frac{b_1}{y-i} + \frac{b'_1}{y+i} + \frac{P'_n(y)}{P_n(y)} + C, \quad (3.46)$$

where b_1 and b'_1 are the residues at $y = i$ and $y = -i$ respectively and $P_n(y)$ is an n^{th} degree polynomial. C represents the analytic part of χ and from (3.45) one can see that χ is bounded for large y . Thus, from Louville's theorem, C is a constant and χ is a rational function of y . The residues at the fixed poles $y = \pm i$, when calculated in the same way as in the previous chapter gives,

$$b_1 = \frac{1 \pm \lambda}{2} ; \quad b'_1 = \frac{1 \pm \lambda}{2}. \quad (3.47)$$

We assume that χ has finite number of singularities in the complex plane, which is equivalent to saying that the point at infinity is an isolated singularity. As mentioned in the end of §2. 4, one can make use of the fact that

for a rational function, the sum of all the residues is equal to zero, to find the eigenvalues, which gives

$$b_1 + b'_1 + n = d_1, \quad (3.48)$$

where d_1 is the residue at infinity and is calculated by taking Laurent expansion of χ around the point at infinity as

$$\chi = d_0 + \frac{d_1}{y} + \frac{d_2}{y^2} + \dots, \quad (3.49)$$

which when substituted in the QHJ (3.45), gives

$$d_1^2 - d_1 + \left(\frac{1}{4} - s^2\right) = 0 \quad (3.50)$$

from which, the values of d_1 turn out to be

$$d_1 = \frac{1 \pm 2s}{2}. \quad (3.51)$$

Substituting the values of the residues in (3.48), one obtains

$$n = -\frac{1}{2} \pm s \mp \lambda, \quad (3.52)$$

which gives the degree of the polynomial $P_n(y)$ in (3.46). From (3.41), one can see that if $E < 0$, λ becomes imaginary. Hence, (3.52) is not satisfied. Thus, from the above equation, we have the condition $E > 0$, which in turn implies $\lambda > 0$ and real. Hence, one can say that for any range of s , the energy eigenvalues are greater than zero. With this condition on λ , we now proceed to select the values of residues at the fixed poles and at infinity, which will give us the physically acceptable results.

3.3.2 Choice of the residues

First, we shall fix the value of the residue at infinity. From the prior discussion of the potential, we know that the wave functions should not become infinite anywhere, in particular, for $x = \pm a$, *i.e.* the wave function $\psi(y)$ should be finite for large y . From (2.74), the wave function in terms of χ becomes

$$\psi(y) = \exp \left(\int \left(\chi - \frac{y}{1+y^2} \right) dy \right). \quad (3.53)$$

For large y , the leading behaviour of χ is obtained as $\chi \sim \frac{d_1}{y}$ from (3.49), which when substituted in (3.53), gives

$$\psi(y) \sim \exp \left(\int \left(\frac{d_1}{y} - \frac{y}{1+y^2} \right) dy \right) \quad (3.54)$$

$$\sim \frac{y^{d_1}}{(y^2 + 1)^{1/2}}. \quad (3.55)$$

Substituting the value of d_1 from (3.51) in the above equation, one obtains

$$\psi(y) \sim \frac{y^{\frac{1}{2} \pm s}}{(y^2 + 1)^{1/2}}. \quad (3.56)$$

For $0 < s < \frac{1}{2}$, from the above equation, one can see that $\psi \rightarrow 0$ for both values of d_1 and this range corresponds to the case, where the potential exhibits band structure.

For $s > 1/2$, $\psi(y) \rightarrow 0$, only if d_1 takes the value $1/2 - s$. Thus, we fix the value of $d_1 = 1/2 - s$. In this way, the two different ranges of the potential parameter s emerge simultaneously fixing the values of d_1 . To select the values of b_1 and b'_1 , we note that the bound state and band edge wave

functions of one dimensional potentials are non- degenerate, which implies, the wave functions are real. Hence, χ must be real, for which we must have

$$b_1 = b'_1. \quad (3.57)$$

With the above constraint, (3.48) becomes

$$2b_1 + n = d_1. \quad (3.58)$$

Thus, we see that the property, the wave function should be finite for $x \rightarrow \infty$, gives the values of d_1 in the two ranges as follows

$$d_1 = \begin{cases} \frac{1+2s}{2} & \text{for } 0 < s < 1/2 \\ \frac{1-2s}{2} & \text{for } s > 1/2. \end{cases} \quad (3.59)$$

From the parity constraint, one obtains the restriction on the values of the residues at the fixed poles as $b_1 = b'_1$. Using these results, we proceed to calculate the solutions for the two ranges.

3.3.3 Case 1 : Band spectrum

In the range $0 < s < 1/2$, we have seen that d_1 can take both the values of the residues. Hence, taking all the possible combinations of the residues, keeping $b_1 = b'_1$ and substituting them in (3.48), we evaluate for n , which gives the degree of the polynomial $P_n(y)$ and takes the integer values greater than or equal to zero. Thus, there will be four combinations forming four different sets with the expression for n , given in the fifth column of table 3.1. Since n needs to be positive, from table 3.1, we pick only those sets which give a positive integer value for n . As seen in §3.3.1, λ is a positive

Table 3.1: All possible combinations of residues for the range $0 < s < 1/2$

Set	b_1	b'_1	d_1	$n = d_1 - b_1 - b'_1$	remark
1	$\frac{1-\lambda}{2}$	$\frac{1-\lambda}{2}$	$d_1 = 1/2 - s$	$\lambda - s - \frac{1}{2}$	$\lambda > (s + \frac{1}{2})$
2	$\frac{1-\lambda}{2}$	$\frac{1-\lambda}{2}$	$d_1 = 1/2 + s$	$\lambda + s - \frac{1}{2}$	$\lambda > -(s - \frac{1}{2})$
3	$\frac{1+\lambda}{2}$	$\frac{1+\lambda}{2}$	$d_1 = 1/2 - s$	$-\lambda - s - \frac{1}{2}$	not valid
4	$\frac{1+\lambda}{2}$	$\frac{1+\lambda}{2}$	$d_1 = 1/2 + s$	$-\lambda + s - \frac{1}{2}$	not valid

real constant. Thus, only the sets 1 and 2 will give positive values for n and hence, the other two sets namely 3 and 4 are ruled out. So, taking the values of b_1, b'_1 and d_1 from the sets 1 and 2 and substituting them in (3.58), and using the definition of λ and s from (3.41), we obtain the expressions for the energy eigenvalues corresponding to the two band edges of the n^{th} band, as

$$E_n^\pm = \frac{\pi^2}{2ma^2} \left(n + \frac{1}{2} \pm s \right)^2. \quad (3.60)$$

Here, $E_n^\pm$ correspond to the upper and lower band energies of the n^{th} band and they match with the solutions given in [4, 5].

In order to obtain the expression for the wave function, we substitute χ from (3.46) in (3.53) with $b_1 = b'_1$, which gives

$$\psi(y) = (y^2 + 1)^{b_1} \frac{P'_n(y)}{P_n(y)}. \quad (3.61)$$

To evaluate the expression for the polynomial, we substitute χ from (3.46), with $b_1 = b'_1$, in the QHJ equation (3.45), which gives the second order

differential equation

$$P_n''(y) + \left(\frac{4b_1 y}{y^2 + 1} \right) P_n'(y) + \left(\frac{1/4 - s^2}{y^2 + 1} + \frac{4b_1^2 y^2 + 2b_1(1 - y^2) + \lambda^2 - 1}{(y^2 + 1)^2} \right) P_n(y) = 0. \quad (3.62)$$

The sets 1 and 2 have $b_1 = (1 - \lambda)/2$, which on substitution in the above equation, gives

$$P_n''(y) + \left(\frac{2(1 - \lambda)y}{y^2 + 1} \right) P_n'(y) + \frac{\frac{1}{4} - s^2 + \lambda^2 - \lambda}{(y^2 + 1)} P_n(y) = 0. \quad (3.63)$$

From (3.41) and (3.60), one can see that λ has two values $\lambda = n \pm s + \frac{1}{2}$. Substituting these in the above equation, one obtains two differential equations corresponding to the two energy eigenvalues $E_n^\pm$ as

$$(y^2 + 1)P_n''(y) + (1 - 2n \mp 2s)yP_n'(y) + n(n \pm 2s)P_n(y) = 0. \quad (3.64)$$

Defining $y = it$, the above equation takes the form of the well known Jacobi differential equation

$$(1 - t^2)P_n''(t) + (\nu_1 - \nu_2 - t(\nu_1 + \nu_2 + 2))P_n'(t) + n(n + \nu_1 + \nu_2 + 1)P_n(t) = 0, \quad (3.65)$$

with $\nu_1 = \nu_2 = -n \mp s - 1/2$, for the corresponding two λ values. Hence, the polynomial $P_n(t)$ will be proportional to the Jacobi Polynomial and the expression for the two band edge wave functions for the n^{th} band will be

$$\psi(y) = (y^2 + 1)^{\frac{\lambda}{2}} P_n^{\nu_1, \nu_2}(-iy), \quad (3.66)$$

with respective ν_1, ν_2 values corresponding to $\lambda = n \pm 2 + 1/2$.

3.3.4 Case 2 : Bound state spectrum

We proceed in the same way as done in case 1 *i.e.* take all possible combinations of b_1, b'_1 and d_1 , keeping $b_1 = b'_1$ in (3.58). Since, d_1 can take only single value $1/2 - s$, only two sets are possible. Out of these, the set corresponding to $b_1 = b'_1 = (1 - \lambda)/2$ alone, will give a positive value for n . Thus, substituting these values of residues in (3.58), one obtains the expression for the energy eigenvalue as

$$E_n = \frac{\pi^2}{2ma^2} \left(\frac{1}{2} + n + \sqrt{\frac{1}{4} - \frac{2mV_0a^2}{\pi^2\hbar^2}} \right)^2, \quad (3.67)$$

where n can take values 0, 1, 2... and is left with the Jacobi differential equation in terms of t as

$$(1 - t^2)P_n''(t) - 2t(-n - s + \frac{1}{2})P_n'(t) - n(n + 2s)P_n(t). \quad (3.68)$$

The expression for the wave function from (3.53) becomes

$$\psi(y) = (y^2 + 1)^{-\frac{\lambda}{2}} P_n^{s_1, s_2}(-iy), \quad (3.69)$$

where $s_1 = s_2 = -n - s - 1/2$.

Note that by subtracting off the ground state energy and redefining $\alpha = -1$, $A = -(s + 1/2)$ and $x \rightarrow \pi x/a$ in

$$V = A(A - \alpha)\operatorname{cosec}^2 \alpha x - A^2, \quad (3.70)$$

the supersymmetric potential [1], one can see that the above expression and its solutions match with those obtained here.

References

1. F. Cooper, A. Khare and U. Sukhatme *Supersymmetry in Quantum Mechanics* (World Scientific, Singapore, 2001) and references therein.
2. F. Cooper, A. Khare and U. Sukhatme, *Phys. Rep.* **251**, 267 (1995).
3. R. S. Bhalla, A. K. Kapoor and P. K. Panigrahi *Int. J. Mod. Phys. A* **12**, No. **10**, 1875 (1997) and references therein.
4. F. L. Scarf, *Phys. Rev.* **112**, 1137 (1958) and references therein.
5. H. Li and D. Kusnezov, *Phys. Rev. Lett.* **83**, 1283 (1999).
6. S. Sree Ranjani, K. G. Geojo, A. K Kapoor and P. K. Panigrahi, to be published in *Mod. Phys. Lett. A* **19**, No. **19**, 1457 (2004); preprint quant - ph/0211168.
7. R. Dutt, A. Khare and U. Sukhatme, *Phys. Lett. A* **174**, 363 (1993).

Chapter 4

Exactly Solvable Periodic Potentials

4.1 Introduction

This chapter is devoted to the study of potentials, which are ES and periodic in nature. It is well known that periodic potentials play a significant role in condensed matter physics. The energy spectrum of periodic potentials is unique due to the existence of energy bands and the solutions of the Schrödinger equation have the Bloch form:

$$\psi(x) = u(x) \exp(ik.x) \quad (4.1)$$

Here, $u(x)$ has the periodicity of the potential function. These features of the periodic potentials are usually illustrated using the Kronig - Penney model [1], in which the crystal lattice is approximated by a periodic array of square wells. Imposing the Bloch condition on the solutions of the Kronig-Penney model, one is led to a transcendental equation, which has to be solved, in order to obtain the band edge solutions.

Recently, there has been considerable interest in the study [2 - 7] and construction [4, 5, 8, 9] of new periodic potentials, especially those belonging

to the elliptic class. The manifestation of periodic lattices in BEC has also renewed interest in this area [10,11]. This gave us enough motivation to check, if we could apply the QHJ formalism to these potentials. In the previous chapter we have discussed the Scarf potential [12, 13], $V(x) = -\frac{a^2(\frac{1}{4}-s^2)}{2m \sin^2(\frac{\pi x}{2})}$, which exhibits band structure in a finite domain $0 < s < 1/2$ and obtained the band edge solutions of this potential.

Here, we study the ES case of Lamé and the associated Lamé potentials, which belong to the class of elliptic potentials and obtain the band edge solutions [14]. These potentials are usually expressed in terms of the Jacobi elliptic functions $\text{sn}(x, m)$, $\text{cn}(x, m)$ and $\text{dn}(x, m)$ [15, 16], where the parameter m is known as the elliptic modulus, whose value lies between 0 and 1.

The Lamé potential

$$V(x) = j(j+1)m \text{sn}^2(x, m), \quad (4.2)$$

with j being an integer, has $2j+1$ bands followed by the continuum. This potential has been extensively studied in [17 - 19]. It is interesting to note that the Schrödinger equation with the Lamé potential,

$$\frac{d^2\psi(x)}{dx^2} + \frac{2m}{\hbar^2}(E - j(j+1)m \text{sn}^2(x, m))\psi(x) = 0 \quad (4.3)$$

is a result of separating the Laplace equation in the ellipsoidal coordinates and is referred to as the Lamé equation [17, 19]. This has unique analytical properties [17], which render it useful in the study of topics ranging from astrophysics [20 - 26], condensed matter physics [27] to chaos [28,29].

In comparison to the Lamé potential, the associated Lamé potential

$$V(x) = pm \operatorname{sn}^2(x, m) + qm \frac{\operatorname{cn}^2(x, m)}{\operatorname{dn}^2(x, m)}, \quad (4.4)$$

with $p = a(a + 1)$ and $q = b(b + 1)$, did not have a systematic study till recently and has a few scattered references in the mathematical literature. This potential is ES, when $a = b = j$ with j being an integer, and QES when $a \neq b$ with a, b being real. A systematic study of the band structure of this potential has been done for various cases of a and b by Khare and Sukhatme [30] and the algebraic aspects of the QES associated Lamé potential have been investigated in [31 - 35].

The band edge eigenfunctions and the eigenvalues of both these potentials have been found using the conventional techniques for solving the ordinary differential equations [17, 19]. One can obtain the solutions, using the same method used to solve the band structure case of the Scarf potential in the previous chapter.

In §4.2, we show, how one can obtain the form of the wave functions for the Lamé potential for j being an integer, followed by §4.3, where we take up a special case of $j = 2$ and obtain explicit band edge wave functions and energies. In §4.4, the form of the wave functions for the general associated Lamé potential (ES case) is derived, followed by the specific example, where of $a = b = 1$, in §4.5.

4.2 General Lamé potential

The QHJ equation in terms of q for the Lamé potential is

$$q^2 + \frac{dq}{dx} + E - j(j+1)m \operatorname{sn}^2(x, m) = 0. \quad (4.5)$$

As discussed in §2.4, we perform a change of variable

$$t = \operatorname{sn}(x, m), \quad (4.6)$$

which gives $F(t) = \sqrt{(1-t^2)(1-mt^2)}$. The elliptic function is periodic with $\operatorname{sn}(x, m) = \operatorname{sn}(x + 4K(m), m)$, where $K(m)$ is $\int_0^1 dt / \sqrt{(1-t^2)(1-mt^2)}$, the complete elliptic integral of the first kind [15, 16, 36]. We would like to point out here, that the Lamé equation can be written in five forms, namely, two algebraic, one trigonometric, one Weierstrassian and one Jacobi [17] depending on the change of variable. Of all these change of variables, we found (4.6), which gives the Jacobi form of this Lamé equation, to be the best suited for our method. It enables one to write the QHJ equation in a form, which can be easily analyzed using the QHJ formalism. Another added advantage of this particular choice is, that it maps half of the period parallelogram $0 \leq x < 2K(m)$ of the Jacobi elliptic $\operatorname{sn}(x, m)$ function to the upper half complex plane [36]. Using the transformation equations (2.70), one gets the equation for χ as

$$\chi^2 + \frac{d\chi}{dt} + \frac{(mt)^2 + 2m}{4(1-mt^2)^2} + \frac{t^2 + 2}{4(1-t^2)^2} + \frac{2E - 2j(j+1)mt^2 - mt^2}{2(1-t^2)(1-mt^2)} = 0 \quad (4.7)$$

with

$$q = \sqrt{(1-t^2)(1-mt^2)}\phi, \quad \phi = \chi + \frac{1}{2} \left(\frac{mt}{1-mt^2} + \frac{t}{1-t^2} \right). \quad (4.8)$$

The following properties of the Jacobi elliptic functions [15]

$$\frac{d}{dx} \operatorname{sn}(x, m) = \operatorname{cn}(x, m) \operatorname{dn}(x, m) \quad (4.9)$$

and

$$\operatorname{sn}^2(x, m) + \operatorname{cn}^2(x, m) = 1, \quad \operatorname{sn}^2(x, m) + m \operatorname{dn}^2(x, m) = 1 \quad (4.10)$$

were used. For all our further calculations we will regard χ as the QMF instead of p . Using (4.7), in place of the original QHJ equation (4.5), one can obtain the expressions for the wave functions by analyzing the singularity structure of χ .

4.2.1 Form of QMF χ

Equation (4.7) shows that χ has fixed poles at $t = \pm 1$ and $t = \pm 1/\sqrt{m}$. We assume, there are a finite number of moving poles in the complex plane. Hence, we make an assumption that the point at ∞ is an isolated singular point and there are no singularities of the QMF except for those mentioned above. Therefore, proceeding in the same way as done in previous chapters, we can write χ , separating it into its singular and analytical parts as

$$\chi = \frac{b_1}{t-1} + \frac{b'_1}{t+1} + \frac{d_1}{t-1/\sqrt{m}} + \frac{d'_1}{t+1/\sqrt{m}} + \frac{P'_n(t)}{P_n(t)} + Q(t). \quad (4.11)$$

Here, $Q(t)$ is the analytic part of χ and the rational terms represent the singular parts. The coefficients b_1 , b'_1 and d_1 , d'_1 are the residues at $t = \pm 1$ and $t = \pm 1/\sqrt{m}$ respectively. The term $P'_n(t)/P_n(t)$ represents the singular part coming from the n moving poles with residue one. $Q(t)$ represents the

analytic part of χ and from (4.7), one can see that χ is bounded at infinity. Hence, from Louville's theorem, $Q(t)$ is a constant.

The residues b_1 , b'_1 , d_1 and d'_1 at the fixed poles can be determined by substituting the Laurent expansion of χ around these poles in (4.7). The two values of the residues at all the fixed poles turn out to be

$$\frac{3}{4}, \quad \frac{1}{4}. \quad (4.12)$$

Note that all the residues are independent of the potential parameter j . Hence, irrespective of the value of j in the potential, the residues at the fixed poles will take only the above values for any Lamé potential.

We need to select the values of the residues, which give rise to acceptable wave functions. Similar to the case of Scarf potential, there is no way of selecting the residues, hence, we accept both the values i.e. $3/4$ and $1/4$ for each of the residues in the present case. The only requirement that restricts possible combinations of values is that of parity, which arises due the fact that the band edge wave functions are non - degenerate. $\psi(t)$ having definite parity implies $\psi(-t) = \pm\psi(t)$, which in turn gives $\chi(-t) = -\chi(t)$. This condition rules out the possibilities $b_1 \neq b'_1$ and $d_1 \neq d'_1$. So, the possible combinations of b_1 , b'_1 , d_1 and d'_1 , which are allowed, are those which satisfy $b_1 = b'_1$ and $d_1 = d'_1$.

We have assumed that χ has finite number of singularities in the complex plane and hence, the point at infinity is an isolated singularity of χ . The form

of (4.7) suggests that χ is bounded at ∞ . Therefore, χ can be expressed as

$$\chi(t) = \lambda_0 + \frac{\lambda_1}{t} + \frac{\lambda_2}{t^2} + \dots \quad (4.13)$$

and the coefficients λ_k 's are fixed using the QHJ equation (4.7) and one gets $\lambda_0 = 0$, and two values for λ_1 as

$$\lambda_1 = j + 1, -j. \quad (4.14)$$

Making use of the fact, that for a rational function, the sum of the residues is equal to zero, we get

$$2b_1 + 2d_1 + n = \lambda_1, \quad (4.15)$$

where, $b_1 = b'_1$ and $d_1 = d'_1$ have been used. As the left hand side is positive, it is clear that the only choice $\lambda_1 = j + 1$ is consistent with the above equation, thus fixing the value of residue at infinity.

Thus from (4.15), one gets the expression for n as

$$n = j + 1 - 2b_1 - 2d_1, \quad (4.16)$$

which gives the degree of the polynomial $P_n(t)$. So making use of all the residue values at the fixed poles keeping the constraint $b_1 = b'_1$ and $d_1 = d'_1$, one obtains four different sets and corresponding four different values of n from (4.16), as listed out in table 4.1

Table 4.1: Number of zeros of the wave function corresponding to the number of moving poles n of χ , form of $\psi(x)$, number of linearly independent solutions and the total number of zeros of $\psi(x)$.

set	b_1	d_1	n	ψ in terms of x	Number of LI solutions		Total zeros of $\psi(x)$ ¹
					$j = 2N + 1$	$j = 2N$	
1	1/4	1/4	j	$P_j(\text{sn } x)$	$N + 1$	$N + 1$	j
2	3/4	1/4	$j - 1$	$\text{cn } x P_{j-1}(\text{sn } x)$	$N + 1$	N	j
3	1/4	3/4	$j - 1$	$\text{dn } x P_{j-1}(\text{sn } x)$	$N + 1$	N	$j - 1$
4	3/4	3/4	$j - 2$	$\text{cn } x \text{dn } x P_{j-2}(\text{sn } x)$	N	N	$j - 1$

¹These constitute of both real and complex zeros.

4.2.2 Explicit forms of the wave functions

One can obtain the form of the wave functions using (2.74). Substituting the relations given in (4.8) in (2.74), one gets the expression for the wave function in terms of χ as

$$\psi(t) = \exp \left(\int \left(\chi(t) + \frac{1}{2} \left(\frac{mt}{1 - mt^2} + \frac{t}{1 - t^2} \right) \right) dt \right). \quad (4.17)$$

Substituting χ from (4.11) gives

$$\psi(t) = \exp \left(\int \left(\frac{(1 - 4b_1)t}{2(1 - t^2)} + \frac{(1 - 4d_1)mt}{2(1 - mt^2)} + \frac{P'_n(t)}{P_n(t)} \right) dt \right), \quad (4.18)$$

which when simplified and written in terms of the original variable x , gives

$$\psi(x) = (\text{cn } x)^\alpha (\text{dn } x)^\beta P_n(\text{sn } x), \quad (4.19)$$

where, $\alpha = \frac{4b_1 - 1}{2}$ and $\beta = \frac{4d_1 - 1}{2}$. For the sake of simplicity, the elliptic modulus m of the Jacobi elliptic functions will be suppressed here onwards. The four different combinations of the residues (given as sets 1 to 4) give rise to four different forms of the band edge wave functions as listed in the table 4.1.

The parity constraint $\chi(-t) = -\chi(t)$ restricts the polynomial $P_n(t)$ to have either only odd or only even powers of $t \equiv \text{sn } x$. In the sixth and the seventh columns of table 4.1, the number of linearly independent solutions, for the two cases, j being odd and j being even, are given in terms of a positive integer N . For odd j , $N = (j - 1)/2$ and for even j , $N = j/2$. In both the cases, the total number of solutions for a particular j is equal to

$2j + 1$. It is easy to see that the four forms and the number of solutions of a particular form obtained here are in agreement with those already known [17]. For a given set of b_1 and d_1 , n is fixed using (4.16) and the differential equation for the unknown polynomial $P_n(t)$ can be obtained by substituting $\chi(t)$ from (4.11) in (4.7), which gives $Q = 0$ and

$$P_n''(t) + 4t \left(\frac{b_1}{t^2 - 1} + \frac{md_1}{mt^2 - 1} \right) P_n'(t) + G(t)P_n(t) = 0, \quad (4.20)$$

where

$$\begin{aligned} G(t) = & \frac{t^2(4b_1^2 - 2b_1 + 1/4) + 1/2 - 2b_1}{(t^2 - 1)^2} \\ & + \frac{(mt)^2(4d_1^2 - 2d_1 + 1/4) + m/2 - 2md_1}{(mt^2 - 1)^2} \\ & + \frac{2E - 2j(j+1)mt^2 + (16mb_1d_1 - 1)mt^2}{2(t^2 - 1)(mt^2 - 1)}. \end{aligned}$$

For each set of residues given in table 4.1, this differential equation is equivalent to a system of n linear equations for the coefficients of different powers of t in $P_n(t)$. The energy eigenvalues are obtained by setting the corresponding determinant equal to zero. We illustrate this process by obtaining the eigenvalues and eigenfunctions explicitly for the case $j = 2$.

4.3 Lamé potential with $j = 2$

Using the procedure described in the previous section, we obtain the eigenvalues and the eigenfunctions for the supersymmetric potential [8]

$$V_-(x) = 6m \operatorname{sn}^2 x - 2m - 2 + 2\delta, \quad (4.21)$$

where $\delta = \sqrt{1 - m + m^2}$. This potential is same as the Lamé potential in (4.2) with $j = 2$, except for an additive constant $2\delta - 2m - 2$, which has been added to make the lowest energy equal to zero. The equation for $\chi(t)$ is

$$\chi^2 + \frac{d\chi}{dt} + \frac{(mt)^2 + 2m}{4(1 - mt^2)^2} + \frac{t^2 + 2}{4(1 - t^2)^2} + \frac{2E + 4(m + 1 - \delta) - 13mt^2}{2(1 - t^2)(1 - mt^2)} = 0. \quad (4.22)$$

From (4.16), the number of moving poles n , for $j = 2$, are

$$n = 3 - 2b_1 - 2d_1, \quad (4.23)$$

where the values of b_1 and d_1 are obtained from and (4.12). For each set of b_1 , d_1 and n , one can write the form and the number of linearly independent solutions by substituting $j = 2$, which gives $N = 1$ in table 4.1. Hence for the case $j = 2$, the form and the number of solutions are as given in table 4.2.

To get the unknown polynomial part $P_n(t)$ of the wave function and the band edge energies, one needs to substitute the different sets of b_1 , d_1 and n from table 4.2 in the differential equation

$$P_n''(t) + 4t \left(\frac{b_1}{t^2 - 1} + \frac{md_1}{mt^2 - 1} \right) P_n'(t) + G(t)P_n(t) = 0, \quad (4.24)$$

where

$$\begin{aligned} G(t) = & \frac{t^2(4b_1^2 - 2b_1 + 1/4) + 1/2 - 2b_1}{(t^2 - 1)^2} \\ & + \frac{(mt)^2(4d_1^2 - 2d_1 + 1/4) + m/2 - 2md_1}{(mt^2 - 1)^2} \\ & + \frac{2E + 4(m + 1 - \delta) + (16b_1d_1 - 13)mt^2}{2(t^2 - 1)(mt^2 - 1)}. \end{aligned}$$

Table 4.2: The form of the wave functions and the number of linearly independent solutions for the Lamé potential with $j = 2$.

set	b_1	d_1	n	ψ in terms of x	Number of LI solutions
1	1/4	1/4	2	$P_2(\text{sn } x)$	2
2	3/4	1/4	1	$\text{cn } x P_1(\text{sn } x)$	1
3	1/4	3/4	1	$\text{dn } x P_1(\text{sn } x)$	1
4	3/4	3/4	0	$\text{cn } x \text{dn } x P_0(\text{sn } x)$	1

Thus, for various sets of residues, one has the band edge energies and the wave functions as:

Set 1 : $b_1 = 1/4$, $d_1 = 1/4$, $n = 2$

Taking $P_2 = At^2 + Bt + C$, the parity constraint implies

$$B = 0. \quad (4.25)$$

Substituting b_1, d_1 and P_2 in (4.24), one gets a 2×2 matrix equation for A and C as follows

$$\begin{pmatrix} E - 2m - 2 - 2\delta & -6m \\ 2 & E + 2m + 2 - 2\delta \end{pmatrix} \begin{pmatrix} A \\ C \end{pmatrix} = 0. \quad (4.26)$$

Equating the determinant of the matrix in (4.26) to zero, one gets the two values for the energy as

$$E_I = 4\delta, \quad E_{II} = 0, \quad (4.27)$$

which in turn give the polynomial as

$$P_I(t) = m + 1 - \delta - 3mt^2, \quad P_{II}(t) = m + 1 + \delta - 3mt^2. \quad (4.28)$$

From (4.19), we see that the band edge wave functions in terms of x will be

$$\psi_I(x) = m + 1 - \delta - 3m \operatorname{sn}^2 x, \quad \psi_{II}(x) = m + 1 + \delta - 3m \operatorname{sn}^2(x). \quad (4.29)$$

Set 2 : $b_1 = 3/4$, $d_1 = 1/4$, $n = 1$

Taking $P_1 = t - t_0$ and substituting in (4.24), one gets $t_0 = 0$ and the band edge eigenvalue as

$$E = 2\delta - m + 2. \quad (4.30)$$

The band edge wave function becomes

$$\psi(x) = \operatorname{cn} x \operatorname{sn} x. \quad (4.31)$$

Set 3 : $b_1 = 1/4$, $d_1 = 3/4$, $n = 1$

Proceeding in the same way as in set 2, one gets

$$E = 2\delta + 2m - 1, \quad \psi(x) = \operatorname{dn} x \operatorname{sn} x. \quad (4.32)$$

Set 4 : $b_1 = 3/4$, $d_1 = 3/4$, $n = 0$

In this case, $P_0(t)$ is a constant and one gets the band edge energy and the wave function to be

$$E = 2\delta - m - 1, \quad \psi(x) = \operatorname{cn} x \operatorname{dn} x. \quad (4.33)$$

Thus, one obtains five band edge wave functions and their corresponding energies, which agree with those given in [8].

4.4 Associated Lamé potential

The associated Lamé potential is ES when $a = b = j$. In this case, the potential expression becomes

$$V(x) = j(j+1)m \left(\operatorname{sn}^2 x + \frac{\operatorname{cn}^2 x}{\operatorname{dn}^2 x} \right). \quad (4.34)$$

The QHJ equation with the associated Lamé potential is

$$q^2 + q' + E - mj(j+1) \left(\operatorname{sn}^2 x + \frac{\operatorname{cn}^2 x}{\operatorname{dn}^2 x} \right) = 0. \quad (4.35)$$

Doing the change of variable $t \equiv \operatorname{sn} x$ and proceeding in the same way as in the case of general Lamé potential described in §4.2, one gets the equation for $\chi(t)$ as,

$$\begin{aligned} \chi^2 + \frac{d\chi}{dt} + \frac{(mt)^2 + 2m(1 - 2j(j+1))}{4(1 - mt^2)^2} + \frac{2 + t^2}{4(1 - t^2)^2} \\ + \frac{2E - mt^2(1 + 2j(j+1))}{2(1 - t^2)(1 - mt^2)} = 0. \end{aligned} \quad (4.36)$$

4.4.1 Form of QMF (χ)

Similar to the Lamé potential, $\chi(t)$ has fixed poles at $t = \pm 1$ and $t = \pm 1/\sqrt{m}$ along with n moving poles in the entire complex t plane. Further, we assume that the point at infinity is an isolated singularity. χ has no other singular points, except at $t = \pm 1$ and $t = \pm 1/\sqrt{m}$. Following the same procedure used in previous chapters, one obtains the residues at $t = \pm 1$ as

$$b_1 = \frac{3}{4}, \frac{1}{4}, \quad b'_1 = \frac{3}{4}, \frac{1}{4}, \quad (4.37)$$

which is independent of j . The residues at $t = \pm 1/\sqrt{m}$ turn out to be

$$d_1 = \frac{3+2j}{4}, \frac{1-2j}{4}, \quad d'_1 = \frac{3+2j}{4}, \frac{1-2j}{4}, \quad (4.38)$$

which are j dependent. Knowing the singularity structure of χ , one can write it as

$$\chi = \frac{b_1}{t-1} + \frac{b'_1}{t+1} + \frac{d_1}{t-1/\sqrt{m}} + \frac{d'_1}{t+1/\sqrt{m}} + \frac{P'_n(t)}{P_n(t)} + C, \quad (4.39)$$

valid for all t similar to the Lamé potential. Assuming that the point at infinity is an isolated singularity, we calculate the residue at infinity by expanding χ in Laurent expansion for large t as

$$\chi = \lambda_0 + \frac{\lambda_1}{t} + \frac{\lambda_2}{t^2} + \dots \quad (4.40)$$

and substituting it in the the QHJ equation (4.36), one obtains the two values for λ_1 as

$$\lambda_1 = j+1, -j. \quad (4.41)$$

For a rational function, the sum of all its residues is equal to zero. Hence, for χ

$$b_1 + b'_1 + d_1 + d'_1 + n = \lambda_1. \quad (4.42)$$

As in the Lamé case, the parity requirement implies $b_1 = b'_1$ and $d_1 = d'_1$ and we get

$$2b_1 + 2d_1 + n = \lambda_1. \quad (4.43)$$

In this case, both the values of λ_1 are allowed because d_1 and d'_1 can take negative values unlike the case of Lamé potential, where $\lambda_1 = -j$ was ruled

out. Thus, one has two cases $\lambda_1 = j + 1$ and $\lambda_1 = -j$. Note that the potential is invariant under the transformation $j \rightarrow -j - 1$ and one has two different values of j , one positive ($j = j' > 0$) and another negative ($j = -j' - 1 < 0$), leading to the same expression for the potential described by (4.4) and hence, the same answers for the eigenfunctions and eigenvalues. Therefore, it is sufficient to restrict j to positive values alone. We consider only the case $\lambda_1 = j + 1$ and neglect the other value $\lambda_1 = -j$. For $\lambda_1 = j + 1$, the values of n , which describe the number of zeros, for different combinations of b_1 and d_1 values, are given in table 4.3.

The sets 3 and 4 give $n = -1$ and $n = -2$ respectively and will not be considered, as n should be greater than or equal to 0. The sets 1 and 2 will give positive values of n , only if j is positive. Since the singularity structure of the associated Lamé potential is same as the Lamé potential, the form of the wave functions will be same as that given in (4.19). As in the Lamé case, the number of linearly independent solutions are given in the table 4.3 in terms of N , for j being odd or even integer.

4.5 Associated Lamé potential with $j = 1$

We perform the calculation with the supersymmetric potential

$$V_-(x) = 2m \operatorname{sn}^2 x + 2m \frac{\operatorname{cn}^2 x}{\operatorname{dn}^2 x} - 2 - m + 2\sqrt{1 - m}, \quad (4.44)$$

corresponding to $j = 1$.

Table 4.3: Values of n and the form of the wave functions, for different combinations of b_1 and d_1 , for $\lambda_1 = j + 1$.

set	b_1	d_1	n	$\psi(x)$ in terms of x	Number of LI solutions		Total zeros of $\psi(x)^2$
					$j = 2N + 1$	$j = 2N$	
1	1/4	$(1 - 2j)/4$	$2j$	$(\text{dn } x)^{-j} P_{2j}(\text{sn } x)$	$2N + 2$	$2N + 1$	$2j$
2	3/4	$(1 - 2j)/4$	$2j - 1$	$\text{cn } x (\text{dn } x)^{-j} P_{2j-1}(\text{sn } x)$	$2N + 1$	$2N$	$2j$
3	1/4	$(3 + 2j)/4$	-1	-	-	-	-
4	3/4	$(3 + 2j)/4$	-2	-	-	-	-

²These constitute of real and complex zeros.

Proceeding in the same way as in the previous sections, the equation for $\chi(t)$ is found to be

$$\begin{aligned} \chi^2 + \frac{d\chi}{dt} &+ \frac{(mt)^2 - 6m}{4(mt^2 - 1)^2} + \frac{t^2 + 2}{4(t^2 - 1)^2} \\ &+ \frac{2E + 4 + 2m - 4\sqrt{1-m} - 5mt^2}{2(1 - mt^2)(1 - t^2)} = 0 \end{aligned} \quad (4.45)$$

and the form of χ is

$$\chi = \frac{b_1}{t-1} + \frac{b'_1}{t+1} + \frac{d_1}{t-1/\sqrt{m}} + \frac{d'_1}{t+1/\sqrt{m}} + \frac{P'_n(t)}{P_n(t)} + C. \quad (4.46)$$

Substituting (4.46) in (4.45), one obtains $C = 0$ and is left with the differential equation

$$P''_n(t) + 4t \left(\frac{b_1}{t^2 - 1} + \frac{md_1}{mt^2 - 1} \right) P'_n(t) + G(t)P_n(t) = 0, \quad (4.47)$$

where

$$\begin{aligned} G(t) = & \frac{t^2(4b_1^2 - 2b_1 + 1/4) + 1/4 - 2b_1}{(t^2 - 1)^2} \\ & + \frac{(mt)^2(4d_1^2 - 2d_1 + 1/4) + -3m/2 - 2md_1}{(mt^2 - 1)^2} \\ & + \frac{2E + 4 + 2m - 4\sqrt{1-m} + (16b_1d_1 - 5)mt^2}{2(t^2 - 1)(mt^2 - 1)}. \end{aligned}$$

Substituting $j = 1$ in table 4.3, one gets the band edge eigenfunctions and eigenvalues from set 1 and set 2 as follows.

Set 1 : $b_1 = 3/4$, $d_1 = -1/4$, $n = 1$

This combination gives only one solution with

$$E = 2 - m + 2\sqrt{1-m}, \quad \psi(x) = \frac{\text{cn } x \text{ sn } x}{\text{dn } x}. \quad (4.48)$$

Set 2 : $b_1 = 1/4$, $d_1 = -1/4$, $n = 2$

This combination gives two solutions. The band edge energies are

$$E_I = 0 , \quad E_{II} = 4\sqrt{1-m} \quad (4.49)$$

and the corresponding wave functions are

$$\psi_I(x) = \frac{1}{m} \left(\operatorname{dn} x + \frac{\sqrt{1-m}}{\operatorname{dn} x} \right) , \quad \psi_{II}(x) = \frac{1}{m} \left(\operatorname{dn} x - \frac{\sqrt{1-m}}{\operatorname{dn} x} \right) , \quad (4.50)$$

which match with the wave functions in [8]. Thus, in this chapter, we have shown that the QHJ formalism successfully yields the band edge eigenvalues and eigenfunctions for ES periodic potentials.

For all the models studied, the parameter n , which appears in the exact quantization condition (2.40) described in chapter II, gives the number of moving poles of the QMF, which in turn correspond to the zeros of the wave function. For ES models [37], these moving poles were found to be confined to the classical region alone. Thus, the n^{th} excited state has n real zeros. The parameter n characterizes the energy level and increasing n increases the energy.

For the QES models, the same n appears as a parameter in the potential and of the possible states, only a few states, which are determined by n , can be obtained analytically. The QMF for all these states will have the same number of moving poles, which are both real and complex. Correspondingly the wavefunctions for each state will have the same number ($\equiv n$) of zeroes (both real and complex), of which the number of real zeros is in accordance

with the oscillation theorem. The number of real zeros increases as one goes from the lowest state to the highest state possible.

On the other hand, the exactly solvable *periodic* potential models show quite a different kind of distribution of the moving poles in the complex plane. Like QES models, the moving poles of the QMF consist of both real and complex poles. But unlike the QES models, where the number of poles of QMF for all the known states is same, for ES periodic potential models one finds groups of solutions with number of moving poles remaining same within a group, but varying from one group to another group of solutions. This point becomes clear from the form of the solutions listed in table 4.1 for the Lamé potential. Different groups of solutions are precisely the different sets of the solutions listed in table 4.1. In each set, the degree of the polynomial $P_n(\text{sn } x)$ is same and the total number of zeros of the band edge wave functions in the interval $0 \leq x < 2K(m)$ are same. For example, for the $N + 1$ linearly independent solutions belonging to set 1, the total number of zeros of the wavefunctions will be j . However, for the solutions belonging to set 2, there will be $j - 1$ zeros of $P_n(\text{sn } x)$ and one zero corresponding to $\text{cn } x$, thus a total of j zeros. For all the sets, the total number of zeros are given in the last column of table 4.1. However, the number of real zeros in a group increases with the increasing energy and correspondingly the number of complex zeros will decrease. For all values of j , we will have utmost four such groups. A similar pattern is observed for ES associated Lamé potential.

References

1. C. Kittel, *Introduction to Solid state Physics*, 191 (Seventh edition, Wiley Eastern Limited, New Delhi, 1995).
2. H. Li and D. Kusnezov, *Phys. Rev. Lett.* **83**, 1283 (1999).
3. G. V. Dunne and M. Shifman, *Annals of Phys.* **299**, 143 (2002).
4. A. Ganguly, *J. Math. Phys.* **43**, 1980 (2002); preprint math - ph/0207028.
5. V. M. Tkachuk and O. Voznyak, *Phys. Lett. A.* **301**, 177 (2002).
6. P. Ivanov, *J. Phys A : Math. Gen.* **34**, 8145 (2001); preprint quant - ph/0008008.
7. O. Rosas - Ortiz, appeared in Proceedings of the IV workshop on *Gravitational and Mathematical Physics*, N. Bretón *et. al* (Eds), Chapala Jalisco, Mexico (2001); preprint math - ph/ 0302189; *Rev. mex. Fis.* **49**, 145 (2003).
8. A. Khare and U. Sukhatme, *J. Math. Phys.* **40**, 5473 (1999); preprint quant - ph/9906044.
9. G. V. Dunne and J. Feinberg, *Phys. Rev.* **D57** 1271 (1998).
10. J. C. Bronski, L. D. Carr, B. Deconinck and J. N. Kutz, *Phys. Rev. Lett.* **86**, 1402 (2001).

11. C. Fort, F. S. Cataliotti, L. Fallani, F. Ferlaino, P. Maddaloni and M. Inguscio, *Phys. Rev. Lett.* **90**, 140405 (2003).
12. F. L. Scarf, *Phys. Rev.* **D57**, 1271 (1998).
13. F. L. Scarf, *Phys. Rev.* **112**, 1137 (1958) .
14. S. Sree Ranjani, A. K. Kapoor and P. K. Panigrahi, to be published in *Mod. Phys. Lett. A* **19**, No. **27**, 2047 (2004); preprint quant - ph/0312041
15. H. Hancock, *Theory of Elliptic Functions*, (Dover Publications, Inc, New York, 1958).
16. I. S. Gradshteyn and I. M. Ryshik, *Table of Integrals, Series and Products*, (Academic Press, 1965).
17. F. M. Arscot, *Periodic Differential Equations* (Pergamon, Oxford, 1964).
18. W. Magnus and S. Winkler, *Hills Equation* (Interscience Publishers, New York, 1966).
19. E. Whittaker and G. N. Watson, *A Course of Modern Analysis* (Cambridge Univ. Press, Cambridge, 1963).
20. F. Finkel, A. González - López, A. L. Maroto and M. A. Rodriguez, *Phys. Rev.* **D62** (2000) 103515; preprint hep - ph/0006117.
21. R. Kantowski and R. C. Thomas; preprint astro - ph/0011176.

22. O. I. Mokhov, preprint math. DG/0201224.
23. M. Bouhmadi - López, L. J. Garay and P. F. González - Diaz, *Phys. Rev.* **D66**, 083504 (2002); preprint gr - qc/0204072.
24. A. V. Razumov and M. V. Saveliev, Published in Proceedings of the International Conference *Selected Topics of Theoretical and Modern Mathematical Physics (SIMI - 96)*, Tbilisi, Georgia (1996); preprint solv - int/9612004.
25. D. J. Fernandez C, B. Mielnik, O. Rosas - Ortiz and B. F. Samsonov, *J. Phys. A.* **35**, 4279 (2002); preprint quant - ph/0303051.
26. I. Bacus, A. Brandhuber and K. Sfetsos, Contribution to the proceedings of the TMR meeting *Quantum Aspects of Gauge Theories, Supersymmetry and Unification*, Paris (1999); preprint hep - th/0002092.
27. M. D. Glinchuk, E. A. Eliseev and V. A. Stephanovich; preprint cond - mat/0103083.
28. M. Brack, M. Mehta and K. Tanaka, *J. Phys. A.* **34**, 8199 (2001); preprint nlin. CD/0105048.
29. M. Brack, S. N. Fedotkin, A. G. Magner and M. Mehta, *J. Phys. A.* **36**, 1095 (2003); preprint nlin. CD/0207043.
30. A. Khare and U. Sukhatme, *J. Math. Phys.* **42** 5652 (2001); preprint quant - ph/0105044.

31. A. V. Turbiner, *Commun. Math. Phys.* **118**, 467 (1988).
32. M. A. Shifman, *Contemp. Math.* **160**, 237 (1998).
33. A. V. Turbiner, *J. Phys. A.* **22** LI (1989).
34. F. Finkel, A. González - López and M. A. Rodriguez, *J. Math. Phys.* **37**, 3954 (1996).
35. Y. Alhasad , F. Gürsey and F. Iachello, *Phys. Rev. Lett.* **50**, 873 (1983).
36. R. V. Churchill and J. W. Brown, *Complex Variables and Applications* (McGraw - Hill Publishing Company, New York, 1990).
37. S. Sree Ranjani, K. G. Geojo, A. K Kapoor and P. K. Panigrahi *Mod. Phys. Lett. A.* **19**, No. **19**, 1457 (2004); preprint quant - ph/0211168.

Chapter 5

Periodic Quasi Exactly Solvable Models

5.1 Introduction

In chapter I, we have mentioned that the QHJ analysis of ordinary QES models *i.e.* potentials with ordinary bound state spectrum, successfully yielded the QES conditions along with the eigenvalues and the eigenfunctions [1,2]. The present chapter is a study of QES periodic potentials. Here, we discuss the QES case of the associated Lamé potential [3]. The QHJ analysis works on certain assumptions on the singularity structure of the QMF, hence, it is essential to see the validity of these assumptions by analyzing as many different models as possible. Therefore, we study the QES associated Lamé potential, even though the method used is very much similar to that used in the previous chapter.

The expression for the associated Lamé potential is

$$V(x) = a(a+1)m\operatorname{sn}^2(x, m) + b(b+1)m\frac{\operatorname{cn}^2(x, m)}{\operatorname{dn}^2(x, m)}. \quad (5.1)$$

Physically, this potential is described as a periodic lattice of period $K(m)$ with the basis composed of two different atoms, which are alternately placed.

For a systematic study of the associated Lamé potential, one is referred to [4, 5]. This potential is ES, when $a = b$ and has been analyzed in the previous chapter, QES properties when $a \neq b$. There are three different QES cases depending on whether a and b are integers or non - integers in [4] and are given below.

Case (i) : For a, b being unequal integers with $a > b > 0$, there will be a bound bands followed by a continuum band. if $a - b$ is odd(even) integer, then there will be b doubly degenerate band edges of period $2K(4K)$ with the corresponding band gaps zero, which cannot be obtained analytically.

Case (ii) : For a, b being half - integers, but $a + b$ and $a - b$ being integers, there will be infinite number of bands with band edge wave functions of period $2K$ or $4K$ depending on whether $a - b$ is an odd or even. Of these infinite bands, there will be $a - b$ bands whose band edges are non-degenerate with period $2K(4K)$ and $b + \frac{1}{2}$ doubly degenerate states of period $2K(4K)$ and can be obtained analytically.

Case (iii) : For a being an integer and b being a half integer or vice versa, one can obtain some exact analytical results for mid - band states.

For all these above mentioned cases, the information regarding the existence of the missing states, mid band states etc. can be acquired by appealing to the oscillation theorem [6, 7].

We proceed with the analysis of the associated Lamé potential with the same change of variable and the transformation equations used in chapter IV. With the same assumptions on the singularity structure of p , we proceed

to obtain the QES condition and the forms of the band edge wave functions for the general associated Lamé potential in the next section. In section §5.3, we analyze case (i), where both a, b are integers taking the values 2 and 1 respectively. Case (ii), where a and b are both half integers with the values $7/2$ and $1/2$ respectively, is analyzed in section §5.4.

5.2 QES condition and the forms of the wave functions

The QHJ equation for the associated Lamé potential, putting $\hbar = 2m = 1$, is

$$p^2 - ip' = \left(E - a(a+1)m \operatorname{sn}^2(x) - b(b+1)m \frac{\operatorname{cn}^2(x)}{\operatorname{dn}^2(x)} \right). \quad (5.2)$$

Note that with the transformation $b \rightarrow -b - 1$ or $a \rightarrow -a - 1$, the potential does not change. Hence, for all our studies, without losing generality, we take a, b positive and $a > b$. Proceeding in the same way as in chapter IV, defining $p \equiv -iq$ and using the change of variable $t \equiv \operatorname{sn}(x)$, one obtains the equation for χ as

$$\chi^2 + \frac{d\chi}{dt} + \frac{m^2 t^2 + 2m(1 - 2b(b+1))}{4(1 - mt^2)^2} + \frac{2 + t^2}{4(1 - t^2)^2} + \frac{2E - mt^2(1 - 2a(a+1))}{2(1 - t^2)(1 - mt^2)} = 0. \quad (5.3)$$

with the help of the transformation equations

$$q = \sqrt{(1 - t^2)(1 - mt^2)}\phi, \quad \phi = \chi + \frac{1}{2} \left(\frac{mt}{1 - mt^2} + \frac{t}{1 - t^2} \right). \quad (5.4)$$

Like all the earlier investigations, we shall treat χ as the QMF and (5.3) as the QHJ equation.

5.2.1 Form of QMF (χ)

From (5.3), we see that χ has fixed poles at $t = \pm 1$ and $t = \pm \frac{1}{\sqrt{m}}$. In addition to the fixed poles, the QMF is assumed to have finite number of moving poles and no other singular points in complex plane. Hence, one writes χ as a sum of the singular and analytical parts as follows

$$\chi = \frac{b_1}{t-1} + \frac{b'_1}{t+1} + \frac{d_1}{t - \frac{1}{\sqrt{m}}} + \frac{d'_1}{t + \frac{1}{\sqrt{m}}} + \frac{P'_n(t)}{P_n(t)} + Q(t), \quad (5.5)$$

where b_1, b'_1 and d_1, d'_1 are the residues at $t = \pm 1$ and $t = \pm \frac{1}{\sqrt{m}}$ respectively, which need to be calculated. $P_n(t)$ is an n^{th} degree polynomial with $\frac{P'_n}{P_n} = \sum_{k=1}^n \frac{1}{t-t_k}$, being the summation of terms coming from the n moving poles with residue one. The function $Q(t)$ is analytic and since χ is bounded at infinity, from Louville's theorem, it is a constant, say C . The residues at the fixed poles can be calculated by taking the Laurent expression around each individual pole and substituting them in (5.3) as in previous chapters. The two values of the residues b_1, b'_1 at $t = \pm 1$ are

$$b_1 = \frac{3}{4}, \frac{1}{4} \quad b'_1 = \frac{3}{4}, \frac{1}{4} \quad (5.6)$$

and at $t = \pm \frac{1}{\sqrt{m}}$, one has the residue values

$$d_1 = \frac{3}{4} + \frac{b}{2}, \frac{1}{4} - \frac{b}{2} \quad d'_1 = \frac{3}{4} + \frac{b}{2}, \frac{1}{4} - \frac{b}{2}. \quad (5.7)$$

As in chapter IV, we consider both the values of the residues. We demand $b_1 = b'_1$ and $d_1 = d'_1$, a condition coming from the parity constraint $\chi(t) = -\chi(t)$. We assume that the point at infinity is an isolated singularity and

hence using the fact, that the sum of all the residues of a rational function is zero and the restriction $b_1 = b'_1$ and $d_1 = d'_1$, we obtain

$$2b_1 + 2d_1 + n = \lambda_1, \quad (5.8)$$

where λ_1 is residue at infinity. Since χ has an isolated singular point at infinity, one can expand χ in Laurent series around the point at infinity as

$$\chi(t) = \lambda_0 + \frac{\lambda_1}{t} + \frac{\lambda_2}{t^2} + \dots \quad (5.9)$$

Substituting (5.9) in (5.3) and comparing the various powers of t , one obtains

$$\lambda_1 = a + 1, \quad -a. \quad (5.10)$$

Taking various combinations of b_1 and d_1 from (5.6) and (5.7), substituting them in (5.8), one obtains the QES condition from each combination as given in table 5.1 for $\lambda = a + 1$. Thus, one sees that all the allowed combinations of residues give one of the forms of QES condition [5], where $n = 0, 1, 2, \dots$. Note that the other value of residue at infinity *i.e.* $\lambda_1 = -a$, when substituted instead of $a + 1$ in (5.8), gives the QES condition for negative values of a, b *i.e.* for $a \rightarrow -a - 1$, $b \rightarrow -b - 1$ in b_1, d_1 .

5.2.2 Forms of wave function

Since the change of variable and the transformation equations are same as in chapter IV, we get the same expression for the wave function as in (4.18) and (4.19) *i.e.*

$$\psi(x) = \exp \left(\int \left(\chi + \frac{1}{2} \left(\frac{mt}{1 - mt^2} + \frac{t}{1 - t^2} \right) \right) dt \right), \quad (5.11)$$

Table 5.1: The quasi - exact solvability condition from the four permitted combinations of b_1 and d_1

set	b_1	d_1	$2b_1 + 2d_1 + n = \lambda_1$	QES condition
1	3/4	$\frac{3}{4} + \frac{b}{2}$	$2 + b + n = a$	$b - a = -n - 2$
2	3/4	$\frac{1}{4} - \frac{b}{2}$	$1 - b + n = a$	$a + b + 1 = n + 2$
3	1/4	$\frac{3}{4} + \frac{b}{2}$	$1 + b + n = a$	$b - a = -n - 1$
4	1/4	$\frac{1}{4} - \frac{b}{2}$	$-b + n = a$	$a + b = n$

which on substitution of χ from (5.5) and written in terms of the original variable x , becomes

$$\psi(x) = (\text{cn } x)^\alpha (\text{dn } x)^\beta P_n(\text{sn } x), \quad (5.12)$$

where $\alpha = \frac{4b_1-1}{2}$, $\beta = \frac{4d_1-1}{2}$. Hence, for each set of b_1, d_1 , one gets a wave function given by (5.12). The degree n of this polynomial, which is obtained from (5.8), as

$$n = a + 1 - 2b_1 - 2d_1, \quad (5.13)$$

which is in terms of either $a + b$ or $a - b$, as evident from table 5.1. The forms of the wave function can be found and are as given in table 5.2 and table 5.3, for the two different cases, when $a + b$ and $a - b$ are odd and even separately.

Case I Both $a + b$, $a - b$ are even : We introduce $N = \frac{a+b}{2}$ and $M = \frac{a-b}{2}$, where M and N are integers, and obtain the forms of the wave functions in table 5.2, in terms of M and N for the four sets of combinations of b_1 and d_1 in table 5.1.

Table 5.2: The form of the wave functions for the four sets of residue combinations when $a + b$ and $a - b$ are even and equal to $2N$ and $2M$ respectively.

set	b_1	d_1	$n = \lambda_1 - 2b_1 - 2d_1$	$n(M, N)$	wave function $\psi(x)$	LI solutions
1	3/4	$\frac{3}{4} + \frac{b}{2}$	$a - b - 2$	$2M - 2$	$\text{cn } x (\text{dn } x)^{1+b} P_{2M-2}(\text{sn } x)$	M
2	3/4	$\frac{1}{4} - \frac{b}{2}$	$a + b - 1$	$2N - 1$	$\frac{\text{cn } x}{(\text{dn } x)^b} P_{2N-1}(\text{sn } x)$	N
3	1/4	$\frac{3}{4} + \frac{b}{2}$	$a - b - 1$	$2M - 1$	$(\text{dn } x)^{b+1} P_{2M-1}(\text{sn } x)$	M
4	1/4	$\frac{1}{4} - \frac{b}{2}$	$a + b$	$2N$	$\frac{P_{2N}(\text{sn } x)}{(\text{dn } x)^b}$	$N + 1$

Case II Both $a + b$, $a - b$ odd : Introducing, $N' = \frac{a+b}{2}$ and $M' = \frac{a-b}{2}$, where M' and N' are integers, we obtain the wave functions in table 5.3, in terms of M' and N' for the four sets of combinations of b_1 and d_1 in table 5.1.

Table 5.3: The form of the wave functions for the four sets of residue combinations when $a + b$ and $a - b$ are odd and equal to $2N' + 1$ and $2M' + 1$ respectively.

set	b_1	d_1	$n = \lambda_1 - 2b_1 - 2d_1$	$n(M, N)$	wave function $\psi(x)$	LI solutions
1	3/4	$\frac{3}{4} + \frac{b}{2}$	$a - b - 2$	$2M' - 1$	$\text{cn } x (\text{dn } x)^{1+b} P_{2M'-1}(\text{sn } x)$	M'
2	3/4	$\frac{1}{4} - \frac{b}{2}$	$a + b - 1$	$2N'$	$\frac{\text{cn } x}{(\text{dn } x)^b} P_{2N'}(\text{sn } x)$	$N' + 1$
3	1/4	$\frac{3}{4} + \frac{b}{2}$	$a - b - 1$	$2M'$	$(\text{dn } x)^{b+1} P_{2M'}(\text{sn } x)$	$M' + 1$
4	1/4	$\frac{1}{4} - \frac{b}{2}$	$a + b$	$2N' + 1$	$\frac{P_{2N'+1}(\text{sn } x)}{(\text{dn } x)^b}$	$N' + 1$

From the forms of the wave functions in tables 5.2 and 5.3, one observes that the number of linearly independent solutions is different for the two cases. The unknown polynomial in the wave function can be obtained by substituting χ from (5.5) in the QHJ equation (5.3), which gives

$$P_n''(t) + 4P_n(t) \left(\frac{b_1 t}{t^2 - 1} + \frac{m d_1 t}{m t^2 - 1} \right) + G(t) P_n(t) = 0 \quad (5.14)$$

where

$$\begin{aligned} G(t) = & \frac{t^2(4b_1^2 - 2b_1 + \frac{1}{4}) - 2b_1 + \frac{1}{2}}{(t^2 - 1)^2} \\ & + \frac{m^2 t^2(4d_1^2 - 2d_1 + \frac{1}{4}) - 2m d_1 + m(\frac{1-2b(b+1)}{2})}{(m t^2 - 1)^2} \\ & + \frac{2E + (16b_1 d_1 - 1 - 2a(a+1))m t^2}{2(1-t^2)(1-m t^2)}. \end{aligned}$$

The above differential equation is equivalent to a system of n linear equations for the coefficients of different powers of t in $P_n(t)$. The energy eigenvalues are obtained by setting the corresponding determinant equal to zero. In the next section, we obtain the band edge wave functions for the associated Lamé potential with $a = 2$, $b = 1$

5.3 Associated Lamé potential with a, b as integers

For this case, we consider the associated Lamé potential with $a = 2$, $b = 1$. Similar to the previous chapter, we work with SUSY potential

$$V_-(x) = 6m \operatorname{sn}^2 x + 2m \frac{\operatorname{cn}^2 x}{\operatorname{dn}^2 x} - 4m. \quad (5.15)$$

This potential is same as (5.1) with $a = 2$ and $b = 1$, except that a constant has been added to make the lowest energy equal to zero. The QHJ equation in terms of χ is

$$\chi^2 + \chi' + \frac{m^2 t^2 - 6m}{4(1 - mt^2)^2} + \frac{2 + t^2}{4(1 - t^2)^2} + \frac{2E + 8m - 13mt^2}{2(1 - t^2)(1 - mt^2)} = 0. \quad (5.16)$$

Apart from n moving poles, χ has poles at $t = \pm 1$ and $t = \pm 1/\sqrt{m}$. As in the previous section, one can write χ with the parity constraint as

$$\chi = \frac{2b_1 t}{t^2 - 1} + \frac{2md_1 t}{t^2 - \frac{1}{m}} + \frac{P'_n(t)}{P_n(t)}, \quad (5.17)$$

which gives the form of χ in the entire complex plane, where $P_n(t)$ is yet to be determined. Note that for this potential the combination $a + b$ and $a - b$ are both odd *i.e.* 3 and 1 respectively. Hence, we use table 5.3 to obtain all the information regarding the residues at the fixed poles, number of moving poles of χ , number of linearly independent solutions and their form for each set etc. by taking the values of $a = 2$, $b = 1$, which give $M' = 0$ and $N' = 1$. The unknown polynomial in the wave function can be obtained from (5.14), where $G(t)$ for this potential turns out to be

$$\begin{aligned} G(t) &= \frac{t^2(4b_1^2 - 2b_1 + \frac{1}{4}) - 2b_1 + \frac{1}{2}}{(t^2 - 1)^2} \\ &+ \frac{m^2 t^2(4d_1^2 - 2d_1 + \frac{1}{4}) - 2md_1 - \frac{3m}{2}}{(mt^2 - 1)^2} \\ &+ \frac{2E + 8m + (16b_1 d_1 - 13)mt^2}{2(1 - t^2)(1 - mt^2)}. \end{aligned} \quad (5.18)$$

Using (5.14) with $G(t)$ from (5.18) and proceeding in the same way as is in the previous section, one gets the explicit expressions for the eigenfunctions and the eigenvalues as given in table 5.4.

Table 5.4: The residues, the value of n , number of linearly independent solutions, the band edge eigenfunctions and eigenvalues are as follows. Here, $a + b = 3$ and $a - b = 1$, which give $N' = 1$ and $M' = 0$.

set	b_1	d_1	n	LI Solution	Eigenfunction $\psi(x)$	Eigenvalues
1	3/4	5/4	-1	-	-	-
2	3/4	-1/4	2	2	$\frac{\text{cn } x}{\text{dn } x}(3m \text{sn}^2 x - 2 \pm \sqrt{4 - 3m})$	$5 - 3m \pm 2\sqrt{4 - 3m}$
3	1/4	5/4	0	1	$\text{dn}^2 x$	0
4	1/4	-1/4	2	2	$\frac{\text{sn } x}{\text{dn } x}(3m \text{sn}^2 x - 2 - m \pm \sqrt{4 - 5m + m^2})$	$5 - 2m \pm 2\sqrt{m^2 - 5m + 4}$

From the table 5. 4, we see that the first set of residues gives $n = -1$, which will not be considered as n cannot be negative. Thus, this particular case of associated Lamé potential has 5 band edge solutions, which can be obtained analytically out of an infinite number of possible states.

5.4 Associated Lamé potential with a and b as half integers

The potential studied here is the supersymmetric associated Lamé potential with $a = 7/2, b = 1/2$, whose expression is

$$V_- = \frac{63}{4}m \operatorname{sn}^2 x + \frac{3}{4}m \frac{\operatorname{cn}^2 x}{\operatorname{dn}^2 x} - 2 - \frac{29}{4}m + \delta_9 \quad (5.19)$$

where $\delta_9 = \sqrt{4 - 4m + 25m^2}$.

$$\chi^2 + \chi' + \frac{2 + t^2}{4(t^2 - 1)^2} + \frac{m^2 t^2 - m}{4(mt^2 - 1)^2} + \frac{4E + 8 + 29m - 4\delta_9 - 65mt^2}{4(t^2 - 1)(mt^2 - 1)} = 0 \quad (5.20)$$

Note that for this case, $a + b = 4$ and $a - b = 3$ are even and odd respectively. Hence, for such cases, one needs to use sets 2 and 4 from table 5.2 and sets 1 and 3 from table 5.3, in order to get the four groups of the eigenfunctions. The solutions for this potential are given in table 5.5.

We see that there is a degeneracy in the band edge energy eigenvalue $14 - 7m + \delta_9$ and all the solutions agree with the known solutions [5]. In this study, we have demonstrated the applicability of QHJ formalism to QES periodic potentials. We have been successful in obtaining the quasi - exact solvability condition and band edge solutions for cases 1 and 2. Case 3, requires some additional work and is not discussed here.

Table 5.5: The residues, the value of n , number of linearly independent solutions, the band edge eigenfunctions and eigenvalues are as follows, with $\delta = \sqrt{4 - 4m + 25m^2}$. Here, $a + b = 4$ and $a - b = 3$, which give $N = 2$ and $M' = 1$.

set	b_1	d_1	n	LI solutions	Eigenfunction $\psi(x)$	Eigenvalues
1	3/4	1	1	1	$\text{cn } x (\text{dn } x)^{3/2} \text{sn } x$	$\delta_9 - m + 2$
2	3/4	0	3	2	$\text{cn } x (\text{dn } x)^{3/2} \text{sn } x$ $\text{cn } x (\text{dn } x)^{-1/2} \text{sn } x (1 - 2\text{sn}^2 x)$	$\delta_9 - m + 2$ $14 - 7m + \delta_9$
3	1/4	1	3	2	$(\text{dn } x)^{3/2} (12m \text{sn}^2 x - 5m - 2 - \delta_9)$ $(\text{dn } x)^{3/2} (12m \text{sn}^2 x - 5m - 2 + \delta_9)$	0 $2\delta_9$
4	1/4	0	4	3	$(\text{dn } x)^{3/2} (12m \text{sn}^2 x - 5m - 2 - \delta_9)$ $(\text{dn } x)^{3/2} (12m \text{sn}^2 x - 5m - 2 + \delta_9)$ $1 - 8\text{sn}^2 x \text{cn}^2 x$	0 $2\delta_9$ $14 - 7m + \delta_9$

Comparing the two different models of periodic potentials, namely the ES and QES, we find that the singularity structure of the QMF in both the cases is similar. For a potential with fixed values of a and b , these four conditions give four groups of solutions as seen as in table 5.2 and 5.3. A comparative analysis of periodic QES models with their non - periodic counterparts, reveals a few differences. For ordinary QES models, one obtains only one QES condition [1] in terms of n , which appears in the quantization condition (2.70). Specifying the value of n , allows one to pick a potential from a family of potentials. Whereas, in the case of the periodic potentials, in general, one obtains four conditions as given in table 5.1. These four conditions can also be viewed as a constraint between a and b . If we select the value of n and the parameter a , one obtains four different values of b . For example, we choose the value $a = 7/2$ and $n = 4$ and substituting these values in the 4 conditions in table 5.1, one obtains the values of b as $-5/2, 3/2, -3/2, 1/2$. Among these, the values $-5/2, 3/2$ correspond to $q = 15/4$ and $-3/2, 1/2$, correspond $q = 3/4$. Thus, we have two potentials corresponding to distinct (p, q) values, namely $(63/4, 15/4)$ and $(63/4, 3/4)$. The form of the solutions for these potentials can be obtained from tables 5.2 and 5.3. Bot these potentials will have a group of levels which have a QMF with four moving poles.

References

1. K. G. Geojo, S. Sree Ranjani and A. K. Kapoor, *J. Phys. A : Math. Gen.* **36**, 4591 (2003); preprint quant - ph/0207036.
2. K. G. Geojo, *Quantum - Hamilton Jacobi Study of Wave functions and energy spectrum of solvable and quasi - exactly solvable models*, Ph. D. thesis submitted to University of Hyderabad (2004).
3. S. Sree Ranjani, A. K. Kapoor and P. K. Panigrahi; preprint quant - ph/0403196.
4. A. Khare and U. Sukhatme, *J. Math. Phys.* **42**, 5652 (2001); preprint quant - ph/0105044.
5. A. Khare and U. Sukhatme, *J. Math. Phys.* **40**, 5473 (1999); preprint quant - ph/9906044.
6. W. Magnus and S. Winkler, *Hill's Equation* (Interscience Publishers, New York, 1966).
7. F. M. Arscot, *Periodic Differential equations* (Pergamon, Oxford, 1964).
8. H. Hancock, *Theory of Elliptic Functions*, (Dover Publications, Inc, New York, 1958).
9. S. Sree Ranjani, A. K. Kapoor and P. K. Panigrahi, *Mod. Phys. Lett. A.* **19**, No. **27**, 2047 (2004); preprint quant - ph/0312041.

Chapter 6

PT symmetric Hamiltonians

6.1 Introduction

Complex Hamiltonians possessing real eigenvalues have attracted considerable attention in the current literature [1 - 9]. These quantal systems are not well understood because of their recent origin and apart from being counter intuitive. These Hamiltonians are characterized by a PT symmetry, which is discrete parity ($x \rightarrow -x$) followed by time reversal ($i \rightarrow -i$) symmetry. In the case, when the wave functions are also PT symmetric, the eigenvalues are real and the violation of PT symmetry by the wave function leads to eigenvalues which are complex conjugate pairs. Apart from identifying new Hamiltonians belonging to this class, the role of various discrete symmetries is also under thorough investigation.

The presence of complex potentials in these systems, makes them ideal candidates to be probed using the QHJ formalism, since this approach has been formulated in the complex domain[10, 11]. It is extremely interesting to investigate the properties of the QMF of the PT symmetric Hamiltonians

and to find out their differences and similarities with ES real potentials and also the QES ones.

In this chapter, we investigate the structure of the QMF of a class of PT symmetric Hamiltonians consisting of ES and QES models, for which the eigenfunctions and the eigenvalues are simultaneously obtained. The differences and similarities of these novel systems, with their ordinary ES and QES counter parts are clearly brought out.

Application of the QHJ formalism to the PT symmetric potentials requires an approach different from that used earlier. In the absence of a generalization of the oscillation theorem, it is not clear whether the quantization rule (2.40) is valid. If so what contour should be used. In this case, the quantization condition still holds for a contour which encloses all the moving poles. We shall assume this to be the case for the PT symmetric models to be taken up in this chapter. Here we will analyze two PT symmetric potentials namely, the Khare - Mandal potential in §6.2 and the complex Scarf potential $V(x) = -A \operatorname{sech}^2 x - iB \tanh x \operatorname{sech} x$, where $A > 0$ in §6.3.

6.2 Khare - Mandal model

The potential expression for the Khare- Mandal model is $V(x) = -(\zeta \cosh 2x - iM)^2$. This potential has either complex or real eigenvalues depending on whether M is odd or even [2, 4]. Using the QHJ formalism, we obtain the QES condition, when M is odd and M is even. We also obtain the explicit expressions for the eigenvalues and eigenfunctions for the cases $M = 3$ and

$M = 2$. The QHJ equation in terms of $q \equiv ip$, keeping $\hbar = 2m = 1$ and after the change of variable $t \equiv \cosh 2x$ becomes

$$q^2 + 2\sqrt{t^2 - 1}\frac{dq}{dt} + E + (\zeta t - iM)^2 = 0. \quad (6.1)$$

The transformation equations in (2.70) for this change of variable become

$$q = 2(\sqrt{t^2 - 1})\phi, \quad \phi = \chi - \frac{t}{2(t^2 - 1)}, \quad (6.2)$$

which transforms (6.1) into

$$\chi^2 + \frac{d\chi}{dt} + \frac{t^2 + 2}{4(t^2 - 1)^2} + \frac{E + (\zeta t - iM)^2}{4(t^2 - 1)} = 0, \quad (6.3)$$

which will be treated as the the QHJ equation and χ as the QMF..

6.2.1 Form of QMF (χ)

We shall assume that χ has a finite number of moving poles in the complex t plane and that the point at infinity is an isolated singularity. Besides the moving poles, χ has fixed poles at $t = \pm 1$. It is seen from (6.3) that the function χ is bounded at $t = \infty$. Assuming that χ has only these above mentioned singularities, we separate the singular part of χ and write it in the following form.

$$\chi = \frac{b_1}{t - 1} + \frac{b'_1}{t + 1} + \frac{P'_n(t)}{P_n(t)} + C, \quad (6.4)$$

where b_1 and b'_1 are the residues at fixed poles $t = \pm 1$ and $P_n(t)$ is a polynomial of degree n . C gives the analytic part of χ and is a constant due to the Louville's theorem. From (6.3), one can see that χ goes as $\pm i\xi/2$, for large t , which are the values of C .

To find the residues at the fixed poles $t = \pm 1$, we expand χ in Laurent series around each pole and proceeding in the same way as in the earlier chapters, one gets the following two values for b_1 and b'_1 as

$$b_1 = \frac{3}{4}, \quad \frac{1}{4}, \quad b'_1 = \frac{3}{4}, \quad \frac{1}{4}. \quad (6.5)$$

It has been assumed that the point at infinity is an isolated singularity. In order to find the residue of χ at infinity, one expands χ in terms of the Laurent series for large t as follows,

$$\chi = a_0 + \frac{\lambda_1}{t} + \frac{\lambda_2}{t^2} + \dots \quad (6.6)$$

and substitution of which in (6.3), gives the residue at ∞ as

$$\lambda_1 = \frac{im\zeta}{4a_0} \quad (6.7)$$

along with $a_0 = \pm \frac{i\zeta}{2}$, due to which the leading behaviour of χ at infinity takes two values, which turns out to be

$$\lambda_1 = \frac{M}{2}, \quad \frac{-M}{2}. \quad (6.8)$$

Hence, using the fact, that for a rational function, the sum of all the residues is equal to zero, one obtains

$$b_1 + b' + n = \lambda_1 \quad (6.9)$$

From (6.5), we see that the right hand side of (6.9) is positive. Hence, for (6.9) to be true, we choose only the positive value of $\lambda_1 = \frac{M}{2}$, which fixes the value of residue at infinity. This means, we choose $C = a_0 = +i\xi/2$, since a_0

gives the leading behaviour of χ at infinity. It should be noted that there is no way of choosing a particular value of residue at the fixed poles, since one does not have information regarding the square integrability of the solutions. Hence, one needs to consider both the values of b_1 and b'_1 . Thus taking all possible combinations of b_1 and b'_1 in (6.9), one obtains the QES condition for each combination along with a constraint on M as given in table 6.1. From

Table 6.1: The QES condition and the number of moving poles of χ for each combination of b_1 and b'_1 .

set	b_1	d_1	$n = \lambda - b_1 - b'_1$	Condition on M	QES condition
1	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{M}{2} - \frac{1}{2}$	$M = \text{odd}, M \geq 1$	$M = 2n + 1$
2	$\frac{3}{4}$	$\frac{3}{4}$	$\frac{M}{2} - \frac{3}{2}$	$M = \text{odd}, M \geq 3$	$M = 2n + 3$
3	$\frac{3}{4}$	$\frac{1}{4}$	$\frac{M}{2} - 1$	$M = \text{even}, M \geq 2$	$M = 2n + 2$
4	$\frac{1}{4}$	$\frac{3}{4}$	$\frac{M}{2} - 1$	$M = \text{even}, M \geq 2$	$M = 2n + 2$

the table 6.1, we see that sets 1 and 2 are valid, only when M is odd and sets 3 and 4 are valid, only when M is even.

6.2.2 Forms of the wave function

Doing the change of variable and writing p in terms of χ in (2.74), one gets the equation for $\psi(t)$ as

$$\psi(t) = \exp \int \left(\frac{b_1}{t-1} + \frac{b'_1}{t+1} + \frac{P'_n(t)}{P_n(t)} + \frac{i\zeta}{2} - \frac{t}{2(t^2-1)} \right) dt. \quad (6.10)$$

Hence, one can substitute the values of b_1 and b'_1 from sets 1 and 2 in (6.10), if M is odd and sets 3 and 4 if M is even, to obtain the form of the wavefunction. The expression for the wave function is in terms of the unknown polynomial $P_n(t)$, where n gives the number of zeros of $P_n(t)$. In order to obtain the polynomial, we substitute χ from (6.4) in (6.3) to get

$$\begin{aligned} & \frac{P_n''(t)}{P_n(t)} + \frac{2P_n'(t)}{P_n(t)} \left(\frac{b_1}{t-1} + \frac{b'_1}{t+1} + \frac{i\zeta}{2} \right) + \frac{b_1^2 - b_1}{(t-1)^2} + \frac{(b'_1)^2 - b'_1}{(t+1)^2} + \frac{t^2 + 2}{4(t^2 - 1)^2} \\ & + \frac{E + (\zeta t - iM)^2 + 8b_1b'_1 - 4\zeta^2(t^2 - 1)}{4(t^2 - 1)} + i\zeta \left(\frac{b_1}{t-1} + \frac{b'_1}{t+1} \right) = 0. \end{aligned} \quad (6.11)$$

This turns out to be equivalent to n linear homogeneous equations, for the coefficients of different powers of t in $P_n(t)$. The energy eigenvalues are obtained by setting the corresponding determinant equal to zero. The explicit eigenvalues and eigenfunctions are obtained for $M = 3$ and $M = 2$ cases.

6.2.3 Case 1 : $M = 3$

Here, M is odd, so we can use sets 1 and 2 from table 6.1 and get the required results as below.

Set 1 : $b_1 = \frac{1}{4}$, $b'_1 = \frac{1}{4}$ and $n = 1$.

This implies P_n is a first degree polynomial, say, $Bt + C$. Substituting these values in (6.11) and comparing various powers of t , one obtains a 2×2 matrix for B and C as follows

$$\begin{pmatrix} 1 + \frac{E-9+\zeta^2}{4} & -i\zeta \\ -i\zeta & \frac{E-9+\zeta^2}{4} \end{pmatrix} \begin{pmatrix} B \\ C \end{pmatrix} = 0. \quad (6.12)$$

Equating the determinant of this matrix to zero, one obtains the two values for energy and the polynomials as

$$E = 7 - \zeta^2 \pm 2\sqrt{1 - 4\zeta^2}, \quad P_1 = \frac{B}{2}(2t - \frac{i}{\zeta}(1 \pm \sqrt{1 - 4\zeta^2})). \quad (6.13)$$

Substituting the values of b_1 , b'_1 and P_1 in (6.10), gives the two eigenfunctions corresponding to the two eigenvalues as

$$\psi(x) = e^{\frac{i\zeta}{2}\cosh 2x} \left(2\cosh 2x - \frac{i}{\zeta}(1 \pm \sqrt{1 - 4\zeta^2}) \right). \quad (6.14)$$

Set 2: $b_1 = \frac{3}{4}$, $b'_1 = \frac{3}{4}$ and $n = 0$.

Here, we see that $n = 0$ implies $P_n(t)$ is a constant. Substituting these values in (6.11) and proceeding in the same as before, one obtains

$$E = 5 - \zeta^2, \quad \psi(x) = e^{\frac{i\zeta}{2}\cosh 2x} \sinh 2x, \quad (6.15)$$

which match with the known results [2, 4].

6.2.4 Case 2 : M=2

In this case, one makes use of sets 3 and 4 in table 6.1 and proceed in the same way as was done for case 1.

Set 3 : $b_1 = \frac{1}{4}$, $b'_1 = \frac{3}{4}$ and $n = 0$.

The eigenvalues and eigenfunctions obtained are

$$E = 3 - \zeta^2 + 2i\zeta, \quad \psi(x) = e^{\frac{i\zeta}{2}\cosh 2x} (\cosh 2x + 1)^{1/2}. \quad (6.16)$$

Similarly, for

Set 4 : $b_1 = \frac{3}{4}$, $b'_1 = \frac{1}{4}$ and $n = 0$,

one obtains

$$E = 3 - \zeta^2 - 2i\zeta, \quad \psi(x) = e^{\frac{i\zeta}{2}\cosh 2x}(\cosh 2x - 1)^{1/2}. \quad (6.17)$$

These match with the solutions given in [4]. Thus, for any given positive value of M , odd or even, one can obtain the eigenvalues and eigenfunctions for the Khare - Mandal potential. In the next section, we go over to the study of complex Scarf -II potential.

6.3 Complex Scarf - II potential

The expression for the potential is

$$V_- = -A \operatorname{sech}^2 x - iB \operatorname{sech} x \tanh x. \quad (6.18)$$

The QHJ equation in terms of q , where $q = \frac{d\psi}{dx}$ is

$$q^2 + \frac{dq}{dx} + E + A \operatorname{sech}^2 x + iB \operatorname{sech} x \tanh x = 0. \quad (6.19)$$

Doing the change of variable $y = i \sinh x$ and proceeding in the same as was done for other the potentials, one obtains the QHJ equation for χ , with $F(y) = i\sqrt{1 - y^2}$, as follows

$$\chi^2 + \frac{d\chi}{dy} + \frac{2 + y^2}{4(1 - y^2)^2} - \frac{E}{1 - y^2} - \frac{A + By}{(1 - y^2)^2} = 0, \quad (6.20)$$

where the transformation equations (2.70) are

$$q = i(\sqrt{1 - y^2})\phi, \quad \chi = \left(\phi - \frac{y}{2(1 - y^2)} \right). \quad (6.21)$$

Along with the n moving poles with residue one, χ has poles at $y = \pm 1$. We assume that except for these poles, there are no other singularities and the point at infinity is an isolated singularity. Proceeding in the same way as in previous section, we obtain the residues at the poles and at infinity as follows. The residue at $y = -1$

$$b'_1 = \frac{1}{2} \pm \frac{1}{2} \sqrt{\frac{1}{4} + A - B}. \quad (6.22)$$

The residue at $y = 1$

$$b_1 = \frac{1}{2} \pm \frac{1}{2} \sqrt{\frac{1}{4} + A + B} \quad (6.23)$$

and the residue at infinity is

$$\lambda = \frac{1}{2} \pm i\sqrt{E}. \quad (6.24)$$

As seen in the earlier section, one can write χ in terms of its analytic and singular parts as

$$\chi = \frac{b_1}{y-1} + \frac{b'_1}{y+1} + \frac{P'_n}{P_n} + C \quad (6.25)$$

where, C is an analytic part of χ and is a constant due to Louville's theorem and it turns out to be zero. Using the fact, that the sum of all the residues is zero for a rational function, we obtain the expression for the energy eigenvalue as

$$-E = (b_1 + b'_1 + n - \frac{1}{2})^2. \quad (6.26)$$

The wave function in terms of χ can be written using (2.74) as

$$\psi(y) = \exp \left(\int \left(\frac{b_1}{y-1} + \frac{b'_1}{y+1} + \frac{P'_n}{P_n} + \frac{y}{2(1-y^2)} \right) dy \right), \quad (6.27)$$

which is equal to

$$\psi(y) = (y-1)^{b_1-\frac{1}{4}}(y+1)^{b'_1-\frac{1}{4}}P_n(y). \quad (6.28)$$

To obtain the polynomial $P_n(y)$, one needs to solve, (6.20) by substituting χ from (6.25), which gives the following differential equation

$$P_n'' + 2P_n' \left(\frac{b_1}{y-1} + \frac{b'_1}{y+1} \right) + G(y)P_n = 0, \quad (6.29)$$

where

$$\begin{aligned} G(y) = & \frac{(4(b_1^2 - b_1 + b'^2 - b'_1)) + 1 + 4E + 8b_1b'_1)y^2}{(y^2 - 1)^2} \\ & + \frac{2y(4(b_1^2 - b_1 - b'^2 + b'_1) - 2B)}{(y^2 - 1)^2} \\ & + \frac{(4(b_1^2 - b_1 + b'^2 - b'_1) + 2 - 4A - 4E - 8b_1b'_1)}{(y^2 - 1)^2}. \end{aligned}$$

Substituting the expression for E from (6.26) in (6.29) gives,

$$(1-y^2)P_n'' + (2(b'_1 - b_1) - 2(b_1 + b'_1)y)P_n' + n(n+2(b_1 + b'_1 - 1) + 1)P_n = 0 \quad (6.30)$$

which is in the form of the Jacobi differential equation and hence the polynomial $P_n(y)$ is proportional to the Jacobi polynomial $P_n^{2b'_1-1, 2b_1-1}(y)$. Thus, the complete expression for the wave function in terms of the residues and the Jacobi polynomial in the x variable becomes

$$\psi(x) = (i \sinh x - 1)^{b_1-\frac{1}{4}}(i \sinh x + 1)^{b'_1-\frac{1}{4}}P_n^{2b'_1-1, 2b_1-1}(i \sinh x). \quad (6.31)$$

Note that in this whole process, we had written the expression of the eigenvalues and the eigenfunctions in terms of b_1 and b'_1 , which have two values. No

particular value has been chosen. Hence, we need to choose one value of each residue to remove this ambiguity. For this case, unlike the Khare-Mandal potential, the solutions are known to be square integrable. Therefore, we make use of square integrability to choose one of the values of residues. We consider two different ranges of the potential parameters A and B and for each range, obtain the solutions.

6.3.1 Case 1 : $|B| > A + \frac{1}{4}$

With this restriction on A and B , the residues at $y = \pm 1$ becomes $b'_1 = \frac{1}{2} \pm \frac{i}{2}\sqrt{B - A - \frac{1}{4}}$ and $b_1 = \frac{1}{2} \pm \frac{1}{2}\sqrt{A + B + \frac{1}{4}}$ and thus, the wave function in (6.31) becomes

$$\psi(x) = (i\sinh x - 1)^{\frac{1}{4} \pm \frac{s}{2}} (i\sinh x + 1)^{\frac{1}{4} \pm \frac{ir}{2}} P_n^{\pm ir, \pm s}(i\sinh x), \quad (6.32)$$

where $r = \sqrt{B - A - \frac{1}{4}}$ and $s = \sqrt{A + B + \frac{1}{4}}$. From the above equation, one can see that for $x \rightarrow \infty$, $\psi(x) \rightarrow 0$, for

$$b'_1 = \frac{1}{2} \pm \frac{i}{2}\sqrt{B - A - \frac{1}{4}} \quad b_1 = \frac{1}{2} - \frac{1}{2}\sqrt{A + B + \frac{1}{4}}, \quad (6.33)$$

with n limited to lie between $0 \leq n < \frac{1}{2}\sqrt{A + B + \frac{1}{4}} - \frac{1}{2}$. Thus, for this parameter range, expression for $\psi(x)$ will be

$$\psi(x) = (i\sinh x - 1)^{\frac{1}{4} - \frac{s}{2}} (i\sinh x + 1)^{\frac{1}{4} \pm \frac{ir}{2}} P_n^{\pm ir, -s}(i\sinh x) \quad (6.34)$$

and the corresponding expression for energy from (6.26) will be

$$E = - \left(n + \frac{1}{2} - \frac{1}{2} \left(\sqrt{A + B + \frac{1}{4}} + \pm i \sqrt{B - A - \frac{1}{4}} \right) \right)^2. \quad (6.35)$$

6.3.2 Case 2 : $|B| \leq A + \frac{1}{4}$

In this case, the wave function in (6.31) is square integrable, if

$$b'_1 = \frac{1}{2} - \frac{1}{2}\sqrt{\frac{1}{4} + A - B}, \quad b_1 = \frac{1}{2} - \frac{1}{2}\sqrt{\frac{1}{4} + A + B}. \quad (6.36)$$

The expression for the wave function is given by,

$$\psi(x) = (i \sinh x - 1)^{\frac{1}{4} - \frac{s}{2}} (i \sinh x + 1)^{\frac{1}{4} - \frac{t}{2}} P_n^{-t, -s}(i \sinh x), \quad (6.37)$$

where $t = \sqrt{\frac{1}{4} + A - B}$ and $s = \sqrt{\frac{1}{4} + A + B}$ and the energy eigenvalue expression in (6.26) becomes

$$E = - \left(n + \frac{1}{2} - \frac{1}{2} \left(\sqrt{\frac{1}{4} + A - B} - \sqrt{\frac{1}{4} + A + B} \right) \right)^2, \quad (6.38)$$

with n restricted by

$$0 \leq n < \frac{1}{2}\sqrt{\frac{1}{4} + A - B} + \frac{1}{2}\sqrt{A + B + \frac{1}{4}} - \frac{1}{2}. \quad (6.39)$$

In conclusion, PT symmetric potentials belonging to the QES and ES class have been investigated through the QHJ formalism. The QES solvable Khare-Mandal potential has complex or real eigenvalues, depending on whether the potential parameter M is odd or even. The singularity structure of the QMF for these two cases is different. For the case, when M is odd, one observes from table 6.1, that the solutions fall into two groups, which consist of solutions coming from sets 1 and 2. For a solution belonging to a particular group, the number of singularities of the QMF are fixed and consists of both real and complex locations. This kind of singularity structure of the

QMF has been observed in the study of periodic potentials [12, 13]. Though the solutions, for M even, fall into two groups coming from sets 3 and 4, they all have the same number of singularities, which again can consist of complex and real poles. This singularity structure is same as that observed in the ordinary QES models [14].

Coming to the case of exactly solvable PT symmetric potential, the location of the moving poles can be either real or complex. In the specific example of complex Scarf potential, it turns out that all the moving poles are off the real line. In contrast for the ordinary ES models [15], the moving poles are always real. For both the cases, the number of moving poles of the QMF characterize the energy eigenvalues.

References

1. M. Znojil, *J. Phys. A : Math. Gen.* **33**, 4561 (2000); preprint quant - ph/9912027.
2. A. Khare and B. P. Mandal; preprint quant - ph/0004019.
3. Z. Ahmed, *Phys.Lett. A.* **282**, 343 (2001).
4. B. Bagchi, S. Mallik, C. Quesne and R. Roychoudhury; preprint quant - ph/0107095.
5. G. Lévai and M. Znojil, *Mod. Phys. Lett. A.* **16**, 1973 (2001); preprint quant - ph/0110064.

6. G. Lévai and M. Znojil, *J. Phys. A* **35**, 8793 (2002); preprint quant - ph/0206013.
7. O. Yesiltas, M. Simsek, R. Sever and C. Tezcan, *Phys. Scripta*, **67**, 472 (2003); preprint hep - ph/0303014.
8. B. Bagchi, C. Quesne and M. Znojil, *Mod. Phys. Lett. A* **16**, 2047 (2001); preprint quant - ph/0108096.
9. B. Bagchi, F. Cannata and C. Quesne, *Phys. Lett. A* **269**, 79 (2000); preprint quant - ph/0003085.
10. R. A. Leacock and M. J. Padgett, *Phys. Rev. Lett.* **50**, 3 (1983).
11. R. A. Leacock and M. J. Padgett, *Phys. Rev.* **D28**, 2491 (1983).
12. S. Sree Ranjani, A. K. Kapoor and P. K. Panigrahi, to be published in *Mod. Phys. Lett. A* **19**, No. **27**, 2047 (2004); preprint quant - ph/0312041
13. S. Sree Ranjani, A. K. Kapoor and P. K. Panigrahi; quant - ph/0403054.
14. K. G. Geogo, S. Sree Ranjani and A. K. Kapoor, *J. Phys. A : Math. Gen.* **36**, 4591 (2003); preprint quant - ph/0207036.
15. S. Sree Ranjani, K. G. Geojo, A. K Kapoor and P. K. Panigrahi, to be published in *Mod. Phys. Lett. A* **19**, No. **19**, 1457 (2004); preprint quant - ph/0211168.

Chapter 7

Conclusions

In this thesis, we have studied various potential models with special spectral properties using QHJ formalism. The various models, being potentials which exhibit different spectra for different ranges of parameters, potentials with band structure and potentials with both real and complex eigenvalues. For all the models studied, we obtained the eigenvalues and the eigenfunctions successfully. Thus, the QHJ formalism provides an alternate method to solve various types of ES and QES models. The added advantage of this method is, that it is very simple and makes use of well known complex variable theorems.

The most important steps in obtaining the solutions, for both ES and QES models, have been, the choice of the change of variable and use of the singularity structure of the QMF in the complex plane. We did not attempt to derive the singularity structure of the QMF within the QHJ formalism. However, we assumed that the singularity structure is very simple *viz. the QMF has a finite number of moving poles*. With this assumption, we could obtain the bound state and band edge solutions in a most straight

forward fashion, even if the equations to be solved were not of a simple form. The fact that one arrived at correct solutions in all the cases, supports a conjecture that the assumption may be true for all ES and QES models. It should be remarked that this assumption is equivalent to the property that ‘the QMF’, becomes a rational function after the suitable transformations. Stated differently, this assumption is equivalent to, the point at infinity being an isolated singular point. In this work our study has been limited to one dimensional models. It will be interesting to investigate if any of these forms of the assumption remain valid for higher dimensional models.

In all the models, the integer n appearing in the exact quantization condition is the number of moving poles of the QMF. For ES models [1] and for Scarf potential, these poles are located on the real axis only, with no moving poles off the real axis. Given a value of n , there is a unique energy level and the corresponding eigenfunction has only real, n , zeros. However, for the QES models [2], n appears as a parameter in the potential and of the infinite possible states, only a finite number, determined by n , of states can be obtained analytically. In the case of QES models, the moving poles appear at complex locations also. Specifying a value of n , selects a potential within a family and does not pick up a unique level for the potential. In fact, all the analytically solvable eigenfunctions have the same number ($\equiv n$) of zeros, of which the number of real zeros are in accordance with the oscillation theorem [3].

The periodic potentials in chapters IV [4], however, show a richer distri-

bution of the moving poles in the complex plane. Like ordinary QES models, the moving poles of the QMF are located both on and off the real axis. Here, in this case, selecting a value of the integer n , picks up a subset of solutions all having n zeros just like QES models. However, unlike the QES models, varying the integer n does not give a new potential, instead it gives a different subset of levels.

The case of QES periodic potentials [5] of chapter V is very similar to the ES periodic potentials, as far as the singularity structure of the QMF is concerned.

The above conclusions have been arrived in the context of specific models studied. However, we expect them to remain generally valid for other one dimensional, periodic and aperiodic, ES and QES potentials. The number of PT symmetric models [6] studied here is too small to allow any similar generalizations.

Our study exhausts analytically solvable bound state and band edge solutions for both ES and QES cases of one dimensional potentials. For problems which are not ES, making a proper guess about the singularity structure of the QMF, does not appear to be possible. A study of the distribution of the moving poles in the complex plane may have a relation to the classical properties of the system. Such a study is interesting in its own right and can be best done numerically. We expect that, here also, the established results in the complex variable theory will provide a useful scheme to obtain numerical, possibly approximate analytical solutions for the wave functions

and energies. Finally, we hope that a useful extension to continuous energy states in one dimension and to non separable models in higher dimensions can be found.

References

1. S. Sree Ranjani, K. G. Geojo, A. K Kapoor and P. K. Panigrahi, to be published in *Mod. Phys. Lett. A* **19**, No. **19**, 1457 (2004); preprint quant - ph/0211168.
2. K. G. Geojo, S. Sree Ranjani and A. K. Kapoor, *J. Phys. A : Math. Gen.* **36**, 4591 (2003); quant - ph/0207036.
3. E. L. Ince, *Ordinary Differential Equations* (Dover Publications, Inc, New York, 1956).
4. S. Sree Ranjani, A. K. Kapoor and P. K. Panigrahi, to be published in *Mod. Phys. Lett. A* **19**, No. **27**, 2047 (2004); preprint quant - ph/0312041
5. S. Sree Ranjani, A. K. Kapoor and P. K. Panigrahi; preprint quant - ph/0403196.
6. S. Sree Ranjani, A. K. Kapoor and P. K. Panigrahi; preprint quant - ph/0403054.